



and Other Interventional Techniques

Laparoscopy for abdominal emergencies

Evidence-based guidelines of the European Association for Endoscopic Surgery

S. Sauerland,^{1*} F. Agresta,² R. Bergamaschi,³ G. Borzellino,⁴ A. Budzynski,⁵ G. Champault,⁶ A. Fingerhut,⁷ A. Isla,⁸ M. Johansson,⁹ P. Lundorff,¹⁰ B. Navez,¹¹ S. Saad,¹² E. A. M. Neugebauer^{1*}

¹ Institute for Research in Operative Medicine, University of Witten/Herdecke, Ostmerheimer Strasse 200, D 51109 Cologne, Germany

² Department of General Surgery, Presidio Ospedaliero di Vittorio Veneto (TV), Via Forlanini 71, 31029 Vittorio Veneto, Italy

³ Department of Research and Development, Forde Health System, Tefre Veg, 6807, Forde, Norway

⁴ I Divisione Clinicizzata di Chirurgia Generale, Ospedale Borgo Trento, Università di Verona, 37126 Verona, Italy

⁵ 2nd Department of General Surgery, Collegium Medicum of Jagiellonian University, ul. Kopernika 21, 31-501 Krakow, Poland

⁶ Department of Digestive Surgery, CHU J. Verdier, Av. du 14, Juillet, 93140, Bondy, France

⁷ Centre Hospitalier de Poissy-Saint Germain, 10, rue du Champ Gaillard, F 78303 Poissy Cedex, France

⁸ Upper Gastrointestinal and Laparoscopic Unit, Ealing and Charing Cross Hospitals, Uxbridge Road, London W5 2HL, United Kingdom

⁹ Department of Surgery, Sahlgrenska University Hospital, S 41345 Gothenburg, Sweden

¹⁰ Department of Obstetrics and Gynaecology, Viborg Sygehus, Heibergs Alle 4, 8800, Viborg, Denmark

¹¹ Service de Chirurgie Generale et Digestive, Hôpital St. Joseph, Rue de la Duchère 6, Charleroi, B 6060 Gilly, Belgium

¹² Department of Abdominal, Vascular and Transplantation Surgery, Merheim Hospital, University of Witten-Herdecke, Ostmerheimer Str 200, 51109 Cologne, Germany

Online publication: 24 October 2005

Abstract

Background: Emergency laparoscopic exploration can be used to identify the causative pathology of acute abdominal pain. Laparoscopic surgery also allows treatment of many intraabdominal disorders. This report was prepared to describe the effectiveness of laparoscopic surgery compared to laparotomy or non-operative treatment.

Methods: A panel of European experts in abdominal and gynecological surgery was assembled and participated in a consensus conference using Delphi methods. The aim was to develop evidence-based recommendations for the most common diseases that may cause acute abdominal pain.

Recommendations: Laparoscopic surgery was found to be clearly superior for patients with a presumable diagnosis of perforated peptic ulcer, acute cholecystitis, appendicitis, or pelvic inflammatory disease. In the emergency setting, laparoscopy is of unclear or limited value if adhesive bowel obstruction, acute diverticulitis, nonbiliary pancreatitis, hernia incarceration, or mesenteric ischemia are suspected. In stable patients with acute abdominal pain, noninvasive diagnostics should

be fully exhausted before considering explorative surgery. However, diagnostic laparoscopy may be useful if no diagnosis can be found by conventional diagnostics. More clinical data are needed on the use of laparoscopy after blunt or penetrating trauma of the abdomen.

Conclusions: Due to diagnostic and therapeutic advantages, laparoscopic surgery is useful for the majority of conditions underlying acute abdominal pain, but non-invasive diagnostic aids should be exhausted first. Depending on symptom severity, laparoscopy should be advocated if routine diagnostic procedures have failed to yield results.

Key words: Abdominal pain — Emergencies — Laparoscopy

Presented at the 13th International Congress of the European Association for Endoscopic Surgery (EAES), Venice, Italy, June 2005

*Conference organizers on behalf of the Scientific Committee of the European Association for Endoscopic Surgery (EAES), c/o EAES Office, Post Office Box 335, 5500 AH Veldhoven, The Netherlands

Correspondence to: E. A. M. Neugebauer

Acute complaints referable to the abdomen are common presentations in surgical emergency departments. Abdominal pain is the leading symptom in this context. In the context of these guidelines, we define acute abdominal pain as any medium or severe abdominal pain with a duration of less than 7 days. Some of the conditions that cause abdominal pain prove to be self-limiting and benign, whereas others are potentially life-threatening. Since it is often difficult to identify patients who have critical problems early in the course of their disease, laparoscopy offers a superior overview of the abdominal cavity with minimal trauma to the patient.

Table 1. Observational studies on the routine use of laparoscopy in unselected patient cohorts

Study year ^a	No. of patients	Percentages of appendicitis/cholecystitis/gynecological disorders	Definitive diagnosis possible (%)	Percentage of laparoscopic/open surgical/conservative therapy	Avoidance of open surgery (%)
Reiertsen et al., 1985 [225]	81	23/0/23	86	0/35/38	38
Paterson-Brown et al., 1986 [211]	125	NA	91	0/30/70	9
Nagy and James, 1989 [193]	31	29/3/23	90	6/45/48	55
Graham et al., 1991 [99]	79	32/NA/35	99	NA/34/NA	66
Schrenk et al., 1994 [236]	15	67/7/7	93	80/20/0	80
Geis and Kim, 1995 [94]	155	66/5/1	99	96/4/0	96
Navez et al., 1995 [198]	225	18/48/5	93	73/27/0	73
Waclawiczek et al., 1997 [282]	172	17/28/NA	NA	65/28/7	72
Chung et al., 1998 [57]	55	22/15/11	100	62/38/0	62
Salky and Edye 1998 [231]	121	50/0/13	98	43/19/38	91
Sözüer et al., 2000 [252]	56	38/4/32	95	64/13/23	87
Ou and Rowbotham 2000 [207]	77	7/1/52	NA	87/12/1	88
Ahmad et al., 2001 [4]	100	37/23/29	NA	81/19/0	81
Lee and Wong, 2002 [157]	137	25/9/39	91	41/16/43	84
Kirshtein et al., 2003 [130]	277	23/1/9	99	75/25/0	75
Sanna et al., 2003 [232]	94	20/6/26	98	88/12/0	88
Agresta et al., 2004 [2]	602	NA/27/61	96	94/16/0	94
Golash and Willson, 2005 [98]	1320	69/1/10	90	83/7/10	93
Majewski, 2005 [176]	108	41/11/15	100	87/13/0	87

NA, not assessed

^a Studies are ordered according to year of publication

On the other hand, the risks of applying laparoscopy to emergency patients include delay to definitive open surgical treatment, missed diagnoses, and procedure-related complications.

Principally, two different clinical scenarios have to be considered. Either a specific condition can be assumed after diagnostic workup or the reason for the abdominal pain has remained uncertain. Therefore, laparoscopy has a diagnostic but also a therapeutic role. The history of diagnostic laparoscopy covers several decades. In an early study from 1975, Sugarbaker et al. [256] showed that in more than 90% of patients a diagnosis can be established by laparoscopy, thereby avoiding non-therapeutic laparotomy in the majority of cases. Table 1 summarizes several cohort studies of diagnostic laparoscopy, which show that over the years increasingly more patients could be successfully managed exclusively by means of laparoscopic surgery. In parallel, specific laparoscopic procedures were evaluated with regard to their effectiveness in the elective and emergency setting. Today, it is possible to hypothesize that all patients with acute abdominal pain would benefit from laparoscopic surgery. It is the aim of these guidelines to define which subgroups of patients should undergo laparoscopic instead of open surgery for abdominal pain.

Methods

Consensus development

In their meeting on September 11th, 2004, the Scientific and Educational Committee of the European Association for Endoscopic Surgery (EAES) decided to focus new clinical guidelines for the role of laparoscopy in abdominal emergencies. These guidelines were primarily intended to supplement the existing guidelines on specific diseases (e.g., appendicitis and diverticulitis) and secondly to define the role of laparoscopy for other, more rare conditions. Based on a review of the

current literature, European experts were invited to participate in the development of the guidelines. All members of the expert panel were asked to define the role of laparoscopy in the various diseases that may underlie abdominal emergencies. For each disease, two experts summarized independently the current state of the art. From these papers and the results of the literature review, a preliminary document with recommendations was compiled.

In April 2005, the expert panel met for 1 day to discuss the text of the guideline recommendations. All key statements were reformulated until a 100% consensus within the group was achieved [190]. Next, these statements were presented to the audience of the annual congress of EAES in June 2005. Comments from the audience were collected and partly included in the manuscript. The final version of the guidelines was approved by all experts in the panel. Each "chapter" consists of a key statement with a grade of recommendation (GoR) followed by a commentary to explain the rationale and evidence behind the statement.

Literature searches and appraisal

We used the Oxford hierarchy for grading clinical studies according to levels of evidence. Literature searches were aimed at finding randomized (i.e., level 1b evidence) or nonrandomized controlled clinical trials (i.e., level 2b evidence). Alternatively, low-level evidence (mainly case series and case reports; i.e., level 4 evidence) was reviewed. Studies containing severe methodological flaws were downgraded. For each intervention, we considered the validity and homogeneity of study results, effect sizes, safety, and economic consequences.

Systematic literature searches were conducted on Medline and the Cochrane Library until June 2005. There were no restrictions regarding the language of publication. Database searches combined the key word laparoscopy (or laparosc* as title word) with a condition-specific keyword (e.g., diverticulitis). We also paid attention to studies that were referenced in systematic reviews or previous guidelines [35, 134, 214, 275].

Results

General remark

The wide variability in experience with laparoscopy makes it necessary to state that the following recom-

Table 2. Randomized and nonrandomized controlled trials comparing laparoscopic and open repair for perforated gastroduodenal ulcers^a

Study, year	LoE	No. of patients	Leakage rates (%)	Total complication rates (%)	Difference in hospital stay (d)
Lau et al., 1996 [153]	1b	48/45	2/2	23/22	±0 n.s. ^b
Siu et al., 2002 [246]	1b	63/58	2/2	25/50	-1 sign. ^c
Johansson et al., 1996 [119]	2b	10/17	10/7	30/20	-1 n.s. ^b
Sø et al., 1996 [250]	2b	15/38	0/0	7/24	-2 n.s. ^b
Miserez et al., 1996 [74, 185]	2b	18/16	NA	50/9	-1 n.s. ^b
Chung et al., 1998 [57]	2b	3/3	NA	NA	-4 sign. ^c
Kok et al., 1999 [135]	2b	13/20	NA	8/15	-1 n.s. ^b
Næsgaard et al., 1999 [191]	2b	25/49	4/0	28/14	±0 n.s. ^b
Bergamaschi et al., 1999 [21]	2b	17/62	0/0	29/34	-2 n.s. ^b
Mehendale et al., 2002 [180]	2b	34/33	0/0	3/6	-5 sign. ^c
Lee et al., 2001 [155]	3b ^d	155/219	13/2	NA	-1 n.s. ^b
Nicolau et al., 2002 [202]	3b ^d	51/105	0/0	6/7	-2 sign. ^c
Seelig et al., 2003 [240]	3b ^d	24/31	4/3	13/26	-2 n.s. ^b
Tsumura et al., 2004 [272]	3b ^d	58/13	NA	5/23	-12 sign. ^c
Lam et al., 2005 [148]	3b ^d	523/1737	NA	3/13	-3 sign. ^c

LoE, level of evidence; NA, not assessed; n.s., not significant

^a Data are shown for laparoscopic / open group. Studies are ordered according to LoE and year of publication

^b Data are difference of medians

^c Data are difference of means

^d Study was downgraded because type of surgery was selected according to the patient's status or because converted cases were not analyzed within the laparoscopic group

mendations are valid only for surgeons or surgical teams with sufficient expertise in laparoscopic surgery.

Gastroduodenal ulcer

If symptoms and diagnostic findings are suggestive of perforated peptic ulcer, diagnostic laparoscopy and laparoscopic repair are recommended (GoR A).

Perforation is the most dangerous complication of gastroduodenal ulcer disease and accounts for approximately 5% of all abdominal emergencies [208, 298]. In perforated peptic ulcer, surgery is generally superior to conservative treatment evidence level (EL) 1b [27, 61]), also because surgical procedures have improved considerably (EL 1a [184]). Laparoscopic repair of perforated ulcer was first reported in 1990 by Mouret et al. [188]. In two randomized trials, laparoscopic surgery was found to be superior to open surgery for perforated ulcers (EL 1b [153, 246]), and other nonrandomized comparison studies are in accordance with these two trials (Table 2). Complication rates in these studies are strongly influenced by the selection of patients for surgery. Contradictory results were found on postoperative pain levels because there appears to be no difference in pain immediately after surgery (when pain is mainly caused by peritoneal inflammation), but laparoscopic patients seemingly experienced less pain later on (when pain is mainly caused by the incision) (EL 2b [21, 135, 185, 191]). Decreased pain may also account for shorter hospital stay and earlier return to normal activities. Long-term results of both procedures showed no major differences in complication or recurrence rates. Mortality was marginally higher after open surgery, although revisional surgery was more frequently required after laparoscopic surgery (EL 2a [152]).

Many patients in these studies received omental patch repair rather than simple suture, but there is nearly no comparative evidence available to decide

which repair technique is superior (EL 2b [44, 137, 155, 178, 194, 247]; EL 4 [44, 137, 178, 194, 247]). One trial by Lau et al. compared patch repair with fibrin sealing without finding any differences (EL 1b [153]). Conversion to an upper midline incision may be necessary in approximately 10–20% of operations, usually for multiple, large, or rear side perforations and for advanced peritonitis (EL 4 [60, 62, 66, 110, 244]). Nevertheless, conversion does not seem to worsen the clinical outcome compared to open surgery (EL 2b [57]). The treatment of bleeding gastroduodenal ulcers was considered to fall outside the field of the current guidelines.

Acute cholecystitis

Patients with acute cholecystitis should undergo laparoscopic cholecystectomy (GoR A). Surgery should be carried out as early as possible after admission (GoR A). In patients unsuitable for early surgery, conservative treatment or percutaneous cholecystostomy should be considered (GoR B).

Laparoscopy is of minor importance in terms of diagnosis of acute cholecystitis. Studies have shown that the following diagnostic criteria define cholecystitis with nearly 100% specificity: (a) acute right upper quadrant tenderness for more than 6 h and ultrasound evidence of acute cholecystitis (the presence of gallstones with a thickened and edematous gallbladder wall, positive Murphy's sign on ultrasound examination, and pericholecystic fluid collections) or (b) acute right upper quadrant tenderness for more than 6 h, an ultrasound image showing the presence of gallstones, and one or more of the following: temperature above 38°C, leukocytosis greater than 10 × 10⁹/L, and/or C-reactive protein level greater than 10 mg/L (EL 1a [270]).

Traditional treatment consisted of open cholecystectomy, which was performed several weeks after an attack or in the acute setting. With the introduction of

Table 3. Randomized and nonrandomized controlled trials comparing laparoscopic and open cholecystectomy for acute cholecystitis^a

Study, year	LoE	No. of patients	Preoperative duration of symptoms	Total complication rates (%)	Difference in hospital stay (d)
Kiviluoto et al., 1998 [131]	1b	32/31	4 d (mean)	3/42	-2 sign. ^b
Johansson et al., 2005 [122]	1b	35/35	72 h (mean)	2/3	-0 sign. ^b
Kum et al., 1994 [144]	2b	66/43	24–96 h	10/9	-1 sign. ^b
Rau et al., 1994 [224]	2b	102/114	NA	9/11	-2 sign. ^c
Carbajo Caballero et al., 1998 [41]	2b	30/30	NA	NA	-7 sign. ^c
Lujan et al., 1998 [170]	2b	114/110	< 72 h	14/23	-5 sign. ^c
Araujo-Teixeira et al., 1999 [12]	2b	100/100	Variable	10/32	-7 sign. ^c
Pessaux et al., 2001 [218]	2b	50/89	NA	18/21	-5 sign. ^c
Chau et al., 2002 [48]	2b	31/42	Surgery performed 2 d (mean) after admission	13/40	-3 sign. ^c
Eldar et al., 1997 [71]	3b ^d	97/146	72 h (median)	17/26	-4 sign. ^b
Glavic et al., 2001 [97]	3b ^d	94/115	72 h (mean)	10/17	-4 sign. ^c
Bove et al., 2004 [33]	3b ^d	87/153	NA	14/NA	NA
Lam et al., 2005 [148]	3b ^d	1223/1408	NA	1/5	-4 sign. ^c

LoE, level of evidence; NA, not assessed

^a Data are shown for laparoscopic/open group. Studies are ordered according to LoE and year of publication

^b Data are difference of medians

^c Data are difference of means

^d Study was downgraded because type of surgery was selected according to the patient's status or because converted cases were not analyzed within the laparoscopic group

laparoscopy for the surgical approach to gallstone disease acute, cholecystitis was initially considered a contraindication. However, with increasing experience, a number of reports became available demonstrating the feasibility of the laparoscopic approach with an acceptable morbidity [143, 144, 286]. Today, there is sufficient evidence to state that laparoscopy is a safe approach, but the question to ask is if it is clearly superior to an open approach. There are several published studies comparing laparoscopic and open cholecystectomy for acute cholecystitis (Table 3). Only two of them are randomized trials (EL 1b [122, 131]). Nearly all comparative studies demonstrated faster recovery and shorter hospital stay in favor of laparoscopy (EL 1a [152]). Similarly, a minilaparotomic cholecystectomy was studied by Assalia et al. (EL 1b [14]), who were able to reduce hospital stay from 4.7 days with open surgery to 3.1 days with minilaparotomy. However, in the most recently published study, the outcome was very similar in the laparoscopic and conventional groups (EL 1b [122]).

The question remains whether the favorable outcome for laparoscopy is a result of altered pathophysiological response to the operation or whether this is due to concomitant changes in postoperative care due to the expected faster recovery from laparoscopic surgery. There is a clear possibility that trials comparing open and laparoscopic procedures contain traditional care regimens that have not been revised in the open treatment groups but have been modified in the laparoscopic groups, thereby favoring, the expected improved outcome after minimally invasive surgery. Several studies in which hospital stay and convalescence were utilized as endpoints may merely reflect traditions of postoperative care and patient expectations associated with open procedures rather than differences between open and laparoscopic surgical techniques. However, even after the advent of fast-track surgery, the existing evidence supports the use of laparoscopy in terms of earlier

postoperative recovery. The basic recommendation should therefore be to offer all patients a laparoscopic approach. If there is no laparoscopically trained surgeon available, the patient should be treated with an open operation in the acute phase of the disease.

The optimal timing of the operation, regardless of whether performed laparoscopically or conventionally, is of major importance. In fact, timing of surgery seems more important than choice of surgical approach. A large number of studies have compared early versus late cholecystectomy for acute cholecystitis (EL 1a [23, 210]; EL 1b [45, 120, 121, 136, 146, 169], EL 2b [24, 25, 49, 69, 93, 102, 133, 139, 173, 199, 215, 220, 242, 258, 273, 285, 295]). However, the time intervals for early, delayed, or interval surgery were inconsistently defined in these studies. It can be concluded from these studies that conversion rates, complication rates, convalescence times, and hospital costs rise in parallel with an increasing delay between admission and operation (EL 5 [96]). Unfortunately, it is impossible to define the exact time limit until which surgery should be performed, but the majority of studies considered a delay of more than 48 or 72 h to be suboptimal. Delaying surgery is considered potentially harmful, especially in patients with a clinical presentation of gangrenous or hemorrhagic cholecystitis (EL 2b [105, 181]), but laparoscopic surgery in these advanced stages of cholecystitis is technically very demanding.

When performing laparoscopic cholecystectomy, the threshold for conversion should be quite low (EL 4 [168]). In many patient series, conversion rates were between 5 and 40% (EL 4 [15, 33, 36, 48, 70, 80, 95, 105, 140, 168, 199, 215, 230, 258, 268, 295])—much higher than in elective cholecystectomy for uncomplicated cholelithiasis. A set of prognostic variables have been identified that predict the need for conversion, such as degree of inflammation, number of previous gallbladder colics, gallstone size, higher age, male gender, obesity, and surgical, expertise (EL 4 [12, 102, 156, 168,

241]). However, these variables do not allow a completely reliable identification of patients in whom laparoscopic cholecystectomy is impossible. Therefore, every surgical procedure for acute cholecystitis should be started laparoscopically, except for patients with general contraindications.

Despite its general superiority, early laparoscopic cholecystectomy may not be possible in all patients. In elderly patients, comorbidities often render early surgery too risky or they simply preclude anesthesia (EL 5 [39]). These cases can only undergo delayed or interval cholecystectomy, although a small study (EL 1b [280]) suggested that a fully conservative treatment can be tried. In the acute phase, percutaneous cholecystostomy has been proposed as a means of alleviating symptoms until definitive treatment can take place (EL 1b [115]; EL 4 [20, 28, 31, 40, 47, 100, 126, 145, 213, 217, 288]). However, one randomized trial from Greece (EL 1b [109]) found that cholecystostomy and conservative treatment performed similarly well, thus justifying the use of both approaches in an individually tailored manner. On the other hand, the benefits of early surgery should not be generally denied to elderly or comorbid patients. With careful anesthesiologic and surgical management, satisfactory results can be achieved in these difficult subgroups (EL 2b [48]; EL 4 [219]).

Acute pancreatitis

Patients with acute biliary pancreatitis should undergo definitive management of gallstones during the same admission (GoR B). After assessment of severity, mild cases should be done within 2 weeks, whereas severe cases should be done when the general condition has significantly improved (GoR C). The bile duct should be imaged to ensure it is clear of stones (intraoperative cholangiography, magnetic resonance cholangiopancreatography, (MRCP), or endoscopic ultrasound) (GoR B).

Acute pancreatitis is a disease entity with manifold etiologies and large differences in clinical appearance but with high morbidity and mortality in more severe cases. Therefore, classification of acute pancreatitis according to severity is crucial for clinical management. Severe disease requires intensive care and CT imaging (EL 5 [195]). Laparoscopy for diagnostic reasons is unnecessary since diagnosis and classification can be based on other criteria and imaging results (EL 5 [34, 65]).

Early pancreatic necrosectomy compared to late or no surgery has been found to be detrimental in various studies (EL 1b [125, 182]; EL 2b[6, 19, 75, 108, 274]). Whenever possible, necrotic tissue should be allowed to demarcate over a few weeks before necrosectomy takes place. Although some situations (e.g. hemorrhage or compartment syndrome) render surgical exploration inevitable, the majority of cases with severe pancreatitis can and should be spared early surgery (EL 1b [167, 237]). If surgery is necessary, minimally invasive techniques can be chosen for exploration, irrigation, necrosectomy, and drainage (EL 2b [91]; EL 4 [107, 209, 297]), but the open approach is still considered the gold standard (EL 4 [195]).

In biliary pancreatitis, two different approaches may be chosen depending on disease severity. In mild biliary pancreatitis, early laparoscopic cholecystectomy with intraoperative cholangiography is the preferred approach (EL 1b [46, 227, 255]; EL 4 [114, 263]; EL 5 [30, 214]). Bile duct clearance is essential to prevent recurrent disease. Therefore, all patients with biliary pancreatitis should undergo definitive treatment at the next best opportunity, preferably during the same hospital admission. There are no studies available to compare a wait-and-see policy versus early removal of bile duct stones, but the risk of a potentially life-threatening recurrent pancreatitis when delaying bile duct clearance is generally considered to be unwarrantable.

There are three different options available to clear the bile duct: endoscopic stone extraction during endoscopic retrograde cholangiopancreatography (ERCP), laparoscopic exploration, and open exploration. Neither the 1998 EAES guidelines on common bile duct stones nor the 2005 UK guidelines on acute pancreatitis, favored one approach over the others (EL 5 [214, 275]). Because the scientific basis for these recommendations is unchanged, all three strategies are still equally recommendable. In general, surgery should only be started after the bile duct has been cleared, unless there is expertise available for intraoperative duct clearance (EL 2b [276]). If MRCP is available for imaging, it allows detection of choledocholithiasis with sensitivity and specificity both over 90% (EL 2a [124]), although the performance of MRCP may be inferior in acute pancreatitis. In most patients, a negative MRCP is sufficient to exclude bile duct stones, thus obviating the necessity of intraoperative clearance (EL 1b [106]). In conclusion, the optimal strategy in most hospitals will depend on the availability of imaging modalities, on the one hand, and surgical expertise with laparoscopic bile duct exploration, on the other hand.

Severe cases of biliary pancreatitis have a high risk of organ failure and death, which usually contraindicates early surgery. Again, bile duct clearance is necessary, but the timing and methods of definitive therapy are different than in mild disease forms. In severe cases, ERCP with or without endoscopic sphincterotomy followed by interval laparoscopic cholecystectomy is common (EL 1a [16]; EL 1b [76, 87, 200, 269], EL 4 [228]; EL 5 [1, 59]). After the publication of several diagnostic accuracy studies with good results (EL 1b [5, 42, 166, 221, 234]), the role of endoscopic ultrasonography (EUS) increased, but the advantage of EUS depends on the prior probability of bile duct stones (EL 2b [13, 229]). As already mentioned, disease classification is the cornerstone of successful therapy (EL 2b [201]). Several different systems have been proposed for defining a presumably severe case of pancreatitis and for describing the clinical course (Ranson score, APACHE II score, inflammatory markers, etc.), but the difficult choice of an optimal system is beyond the scope of these recommendations. The UK guidelines recommend delaying surgery “until signs of lung injury and systemic disturbance have resolved,” which aptly describes the subjective nature of this decision on timing.

Acute appendicitis

Patients with symptoms and diagnostic findings suggestive of acute appendicitis should undergo diagnostic laparoscopy (GoR A) and, if the diagnosis is confirmed, laparoscopic appendectomy (GoR A). If diagnostic laparoscopy shows that symptoms cannot be ascribed to appendicitis, the appendix may be left in situ (GoR B).

Appendicitis is a very common disease, but its symptoms are often equivocal and many other causative pathologies can be responsible. Despite improved imaging with sonography or CT, the rates of false-negative appendectomy are still high, especially in women (EL 4 [29, 86]). Among the 56 randomized trials that have compared laparoscopic and conventional approaches for suspected appendicitis (EL 1a [233]; EL 1b [186]), only a few studies have explicitly used the findings of diagnostic laparoscopy to guide further surgical therapy. Most of these studies included only female patients of fertile age and documented a large reduction in the rate of negative appendectomy (EL 1b [37, 117, 147, 151, 205, 277]). However, the diagnostic advantages in men and children seem to be smaller and less consistent since appendicitis is much easier to diagnose in these subgroups.

The relative advantage of laparoscopic over conventional appendectomy has been under debate for more than a decade. According to the most recent Cochrane Review (EL 1a [233]), laparoscopic appendectomy offers certain advantages, although the difference compared to open appendectomy is not major. The EAES guidelines on appendectomy clearly favor the laparoscopic approach (EL 5 [72]), mainly because of the significantly reduced risk of wound infection and the faster postoperative recovery. This recommendation also pertains to perforated cases.

If the appendix looks normal on laparoscopy but another pathology is found to be the cause of the patient's symptom, then the appendix should be left in situ (EL 4 [278]). The 10-year follow-up by van Dalen et al. (EL 1b [277]) demonstrated the safety of this approach in women. The situation is more complicated when the appendix shows no signs of inflammation and no other pathology can be found. Different groups have provided contradictory data on the reliability of macroscopic diagnosis of appendicitis (EL 4 [51, 103, 141, 266]). Weighing the disadvantage of a negative appendectomy against the risk of overlooking a case of appendicitis is difficult. If symptoms and signs are severe and typical for appendicitis, most surgeons will consider appendectomy to be indicated because in early appendicitis inflammation may be limited to intramural layers.

Acute diverticulitis

Patients with presumed acute uncomplicated diverticulitis should not undergo emergency laparoscopic surgery (GoR C). Although colonic resection remains standard treatment for perforated diverticulitis, laparoscopic lavage and drainage may be considered in some selected patients (GoR C).

After physical examination and a blood count, CT is especially useful to diagnose diverticulitis. If compli-

cated disease is likely, CT is able to visualize inflammation of the pericolic fat, thickening of the bowel wall, or peridiverticular abscess. Diagnostic laparoscopy is therefore unnecessary. Resection of the diseased segment should be performed in an elective rather than an emergency setting since the risk of conversion and the rate of primary reanastomosis strongly depend on the presence and severity of acute inflammation. The value of elective laparoscopic sigmoid resection has been addressed in guidelines issued by the EAES in 1999 [134].

Complicated cases of diverticular disease are classified according to the modified Hinchey classification. Stage I indicates the presence of a pericolic abscess, stage IIa indicates distant abscess amenable to percutaneous drainage, and stage IIb indicates complex abscess associated with or without fistula. Diffuse peritonitis is classified as stage III (purulent) or IV (fecal). Peritonitis or pneumoperitoneum usually require emergency surgical exploration (EL 1b [142, 294]; EL 5 (10, 212)). In Hinchey stages III and IV, laparoscopic abdominal exploration and peritoneal lavage have been successfully used, but there are only limited data available (EL 2b [77]; EL 4 [88, 206, 223, 235]). A laparoscopic approach may be especially advantageous in high-risk patients, who would probably not survive Hartmann's procedure. In such patients, perforation may be closed by an omental patch (EL 4 [88]). In stage IIb, abscesses can be drained and fistula can be closed laparoscopically (EL 4 [88, 223, 238]), but it must be taken into account that only very few surgeons are experienced enough to perform these operations. It is therefore too early to generally recommend laparoscopic emergency surgery for complicated diverticular disease, despite promising results.

Small bowel obstruction due to adhesions

In the case of clinical and radiological evidence of small bowel obstruction nonresponding to conservative management, laparoscopy may be performed using an open access technique (GoR C). If adhesions are found at laparoscopy, cautious laparoscopic adhesiolysis can be attempted for release of small bowel obstruction (GoR C).

The clinical value and the potential complications of adhesiolysis are highly debated. A blinded trial by Swank et al. found similar levels of pain after diagnostic laparoscopy with or without adhesiolysis (EL 1b [262]). Although this trial was performed in patients with chronic recurrent abdominal pain, it also has implications for the acute pain situation. On the other hand, laparoscopic adhesiolysis is sometimes performed at diagnostic laparoscopy for acute abdominal pain, to enable complete visualization of the abdominal content. Therefore, the term adhesiolysis covers a wide spectrum of invasiveness. Furthermore, the natural variability of adhesions and their sequelae determines possible success and failure rates of adhesiolysis. Therefore, the decision for adhesiolysis in the acute setting is a balance of these factors (EL 2b [284]). As a rule, adhesiolysis in an abdomen without intestinal obstruction should be kept to a minimum.

Radiographically confirmed small bowel obstruction requires emergency surgery (EL 2b [82–84] when nonoperative therapy is unsuccessful. Laparoscopic treatment of acute small bowel obstruction was first described by Bastug et al. (EL 4 [18]) and has since been reported by others (EL 4 [3, 8, 17, 32, 55, 56, 89, 90, 111, 129, 160, 163, 197, 243, 254, 259, 261]). Studies comparing the results of laparoscopic and conventional treatment of this condition are nearly lacking, except for the matched-pair analysis by Wullstein and Gross (EL 2b [289]). The benefits of the laparoscopic approach that have been reported consist of a more rapid postoperative recovery with faster return of bowel movements, lower morbidity, and shorter hospital stay. However, there is concern that laparoscopic treatment of small bowel obstruction may lead to a higher rate of bowel injury than conventional surgery. In the single comparative study (EL 2b [289]), the risk of perforation was clearly higher in the laparoscopic group (27%). The high conversion rate is also an issue. Complete laparoscopic treatment seems to be possible in only 50%–60% of patients (EL 4 [3, 8, 17, 32, 55, 56, 89, 111, 129, 160, 163, 197, 243, 254, 259, 261]). The remaining patients have to be converted to open surgery for malignant disease, iatrogenic bowel perforation, or other reasons. Some studies have examined predictive factors for successful laparoscopy (EL 2b [163, 259]). A history of two or more surgical abdominal operations, late operation (> 24 h), and bowel diameter exceeding 4 cm have been reported to be predictors of conversion. An isolated scar from a previous appendectomy seems to be favorable in terms of avoiding a conversion. To avoid the possibility of intraabdominal injuries during laparoscopic access, open rather than laparoscopic surgery should be performed if scars or other findings indicate the presence of severe or extended adhesions (EL 4 [85, 192]).

Incarcerated hernia

Although the open approach remains standard treatment for incarcerated hernia, laparoscopic surgery may be considered in carefully selected patients (GoR C).

The available evidence for the use of laparoscopic surgery in inguinal, incisional, and other hernias is very good, but all these studies have excluded symptomatic and emergency surgery cases. It seems unjustified to adopt the principle of transferable evidence to delineate the treatment of incarcerated hernia from the results obtained in the elective setting. With regard to the laparoscopic treatment of incarcerated hernias, so far only case reports (EL 4 [38, 58, 78, 123, 150, 164, 271, 283, 290]) and small case series (EL 4 [81, 113, 149, 154, 165, 239]) have been published. The largest series is from Leibl et al. (EL 4 [158]) and reports on 220 patients. The authors—highly experienced laparoscopic surgeons—found their results in incarcerated groin hernias to be similar to those elective for hernia repair. Because there are no comparative studies available to compare open and laparoscopic surgery, one should be very reluctant to choose a laparoscopic approach to hernia sac, abdominal wall, or peritoneum. Although early

clinical results are promising, these techniques should be restricted to surgeons with maximum expertise in laparoscopic hernia surgery.

Mesenteric ischemia

If mesenteric ischemia is clinically suspected, conventional imaging is preferable over diagnostic laparoscopy in defining therapeutic management (GoR C).

Acute mesenteric ischemia is caused by arterial occlusion (approximately 50% of cases), nonocclusive arterial ischemia (20–30%), or venous occlusion (5–15%) [253]. A clinical diagnosis of mesenteric ischemia is usually confirmed by the use of conventional angiography, CT scanning, or duplex sonography [132, 204, 216]. Traditional surgical therapy consists of resection of infarcted bowel segments or embolectomy, depending on duration and extent of ischemia. The benefit of surgery needs to be considered on a case-by-case basis since there is no good evidence available to compare surgical and medical treatment for those patients with a salvageable condition (EL 2b [26, 68, 296]).

The potential value of emergency laparoscopy in these patients relates to its diagnostic rather than its therapeutic opportunities. However, the rate of mesenteric ischemia among patients with acute abdomen is only approximately 1% [112]. Furthermore, laparoscopic viewing does not guarantee correct recognition of ischemia. Since radiographic imaging accurately identifies most cases of mesenteric ischemia, it is very unlikely that diagnostic laparoscopy will prevent a negative laparotomy in these patients. In the literature, only a few cases have been published (EL 4 [52, 54, 73, 292]; EL 5 [159]), although there are more reports concerning second-look laparoscopies.

Gynecologic disorders

If gynecologic disorders are the suspected cause of abdominal pain, diagnostic laparoscopy should follow conventional diagnostic investigations (GoR A), and, if needed, a laparoscopic therapy for the disease should be performed (GoR A). A close cooperation with the gynecologist is strongly recommended.

Many acute gynecologic disorders can be approached safely and effectively by laparoscopy with the intent not only to correctly diagnose the patient but also to render treatment (EL 4 [138, 174, 196, 207]). The most common diagnoses encountered in women with acute pelvic pain are ectopic pregnancy (approximately 20% of cases), salpingo-oophoritis (20%), pelvic adhesions (20%), endometriosis (15%), and ovarian cysts (15%). In gynecological emergencies, CT scans are very seldom helpful. After a pregnancy test, transvaginal and conventional ultrasound can aid in formulating a differential diagnosis. However, diagnostic laparoscopy is superior to other diagnostic tools (EL 2b [183]) and may lead to the correction of an erroneous preoperative diagnosis in up to 40% of patients (EL 4 [7, 67, 138, 264]).

Ectopic pregnancy (EP) is a life-threatening condition. In early pregnant women presenting with acute pelvic pain and/or vaginal bleeding, a diagnostic laparoscopy should always be considered to exclude EP. In the vast majority of cases, a pregnancy test can exclude the diagnosis in cases with only minor symptoms. When serum human chorionic gonadotropin (hCG) levels reach 1,000 IU/L, transvaginal ultrasonography can differentiate between an EP or an intrauterine pregnancy (IUP) because all IUPs can clearly be seen in cases with hCG > 1,000 IU/L. A normal IUP will have a hCG doubling rate of 2 days. Thus, vaginal ultrasound and hCG go hand in hand in the diagnosis of EP in cases of minor or no abdominal symptoms (EL 5 [222]). In cases with EP, laparoscopic surgery should be undertaken also because of its total cost is cheaper (EL 1b [101]). It is fast, and fertility outcome is comparable to laparotomy. Furthermore, sick leave and hospitalization are shorter and adhesion development is minor compared to laparotomy (EL 1b [171, 172, 279]; EL 2b [79, 189]). Laparoscopic salpingectomy should be performed in cases of ruptured tubal pregnancy. In cases of unruptured tubal pregnancy, a tube-preserving operation should be considered. Hemodynamic instability is a contraindication for laparoscopy.

Torsion of ovarian cysts is an organs-threatening disease. Patients often present with acute abdominal pain. After excluded pregnancy, a transvaginal ultrasound is mandatory to exclude ovarian cyst formation. In the majority of patients, free fluid can be seen in the abdomen, and if symptoms decline, an expectative attitude can be undertaken. In cases with persistent pain and/or if a larger cyst is seen on ultrasound, a diagnostic laparoscopy must be performed to exclude adnexal torsion. Ovarian cysts that are found during diagnostic laparoscopic should be treated laparoscopically (EL 1b [175, 291]). Pregnant women with acute pelvic pain and clinical signs of torsion of ovarian cyst should be offered laparoscopic repair. Laparoscopic surgery was also reported to be superior compared to open surgery for resecting other types of ovarian cysts (EL 1b [203]).

Endometriosis often causes infertility and pain. Pain is usually chronic and recurrent, but some patients present with acute symptoms. Surgical treatment may be indicated in some patients and may be performed as an open procedure or laparoscopically. Only one trial has compared the two approaches (EL 1b [175]) and documented a significantly faster and less painful recovery after laparoscopy. More evidence is available on the comparative effectiveness of laparoscopic excision versus conservative treatment of endometriosis. Although these studies included elective rather than emergency patients, their results indicate that laparoscopic excision results in clear and patient-relevant advantages as opposed to conservative treatment (EL 1a [116]; EL 1b [1, 260]).

Salpingo-oophoritis commonly causes acute pelvic pain and often mimics other diseases. Conservative treatment consists of antibiotics. Laparoscopy is useful to exclude other pathologies, which may be present in approximately 20% of patients (EL 4 [22]). Furthermore, microbiological specimens can be taken to guide antibiotic therapy. Depending on the severity of symp-

toms, laparoscopy is therefore considered to be advantageous for acute salpingitis (EL 4 [22, 251]) and pyosalpinx (EL 4 [267]).

Nonspecific abdominal pain

Patients with severe nonspecific abdominal pain (NSAP) after full conventional investigations should undergo diagnostic laparoscopy if symptoms persist (GoR A). Patients with NSAP of medium severity may undergo diagnostic laparoscopy after a period of observation (GoR C).

According to symptoms and diagnostic findings, most patients with acute abdominal pain can easily be categorized into different groups of presumed diagnoses, but some patients will not fit into these diagnostic categories due to unclear or equivocal findings. In these cases, of NSAP, the severity of symptoms determines the necessity of emergency surgery. Some patients definitely require surgical exploration, a second group can safely be monitored under conservative therapy, and in a third group the decision between operative or conservative management is unclear. If symptoms are severe enough to require surgical exploration, this should be done laparoscopically. The reason lies more in the therapeutic than the diagnostic value of laparoscopic surgery. As described previously, laparoscopic surgery is advantageous for many intraabdominal diseases, which may also turn out to be the underlying cause of an unclear abdomen. Also, because converted cases have a similar outcome compared to primarily open cases (EL 2b [57]), the benefits of a laparoscopic approach outweigh its potential negative effects.

Four randomized controlled trials have compared early laparoscopy versus observation for nonspecific acute abdominal pain (Table 4). Three trials focused exclusively on right iliac fossa pain in women after excluding clear cases of appendicitis (EL 1b [43, 92, 187]). The fourth trial included 120 men and women with acute abdominal pain regardless of pain localization (EL 1b [64]). Three out of four trials found that early laparoscopy clearly facilitated the establishment of a diagnosis with subsequent therapy, whereas more patients in the control group left the hospital without a clear diagnosis. More important, hospital stay was shorter in two of the trials (EL 1b [43, 92]). At 1-year follow-up, recurrent pain episodes were less frequent (EL 1b [187]) and health-related quality of life was better (EL 1b [64]) in the laparoscopic group. Based on these data, it seems justified to lower the threshold for surgical exploration when using a laparoscopic rather than an open approach. However, it seems advisable to observe patients over some hours because abdominal symptoms may become more specific over time or simply disappear in some cases (EL 4 [128]).

Abdominal trauma

For suspected penetrating trauma, diagnostic laparoscopy is a useful tool to assess the integrity of the peritoneum and avoid a nontherapeutic laparotomy in stable patients

Table 4. Randomized controlled trials comparing laparoscopic surgery and conservative management for acute nonspecific abdominal pain^a

Study, year	LoE	No. of patients	Patients in conservative group receiving surgical exploration (%)	Patients remaining without a final diagnosis (%)	Difference in hospital stay (d)
Champault et al., 1993 [43]	1b	33/32	50	3/72	-2 sign. ^b
Decadt et al., 1999 [64]	1b	59/61	28	19/64	±0 n.s. ^c
Gaitán et al., 2002 [92]	1b	55/55	40	5/2	-1 sign. ^b
Morino et al., 2003 [187]	1b ^d	24/29	31	12/55	NA

LoE, level of evidence; NA, not assessed; n.s., not significant

^a Data are shown for laparoscopic/conservative group. Studies are ordered according to year of publication

^b Data are difference of means

^c Data are difference of medians

^d Only published abstract available

(GoR B). Stable patients with blunt abdominal trauma may undergo diagnostic laparoscopy to exclude relevant injury (GoR C).

Laparotomy for abdominal trauma used to be negative or nontherapeutic in approximately one-third of patients (EL 4 [162, 226]), but modern imaging techniques have reduced this figure to less than 10% (EL 4 [104]). The literature contains approximately 40 prospective or retrospective cohort studies on the diagnostic role of laparoscopy in trauma (EL 4 [281]). The major advantage of laparoscopy as identified in these studies was the obviation of unnecessary laparotomy in approximately 60% of cases. However, relevant injuries went undetected in 1% of all laparoscopies, particularly after blunt trauma affecting solid organs or hollow viscus (EL 4 [281]). Because the majority of the available evidence derives from patients with stab or gunshot wounds, diagnostic laparoscopy seems to be recommendable as a screening tool for patients with a moderate to high index of suspicion for intraabdominal injuries. However, in hemodynamically unstable patients, emergency surgical exploration of the abdomen may be life-saving. In this situation, delaying definitive therapy by laparoscopy is contraindicated.

Two randomized studies have been published on laparoscopy in trauma. A small study compared laparoscopy with peritoneal lavage and found higher diagnostic specificity in the laparoscopic group (EL 1b [63]). The second trial was, in fact, a double trial (EL 1b [161]). First, it compared exploratory laparotomy and diagnostic laparoscopy for stab wounds that had penetrated the peritoneum. Second, patients with equivocal peritoneal violation were randomized to diagnostic laparoscopy or expectant nonoperative management. Not unexpectedly, laparoscopy reduced hospital stay compared to laparotomy but prolonged hospital stay compared to conservative management (EL 1b [161]). Although laparoscopy saved more than half of patients from laparotomy, the postoperative clinical course and costs failed to differ between laparoscopic and laparotomic group. Because the study was relatively small and did not report on the potential long-term advantages of laparoscopy, further research is needed. Accordingly, the panel believes that the available evidence does not justify a high-grade recommendation.

Although the trials mentioned previously did not use laparoscopy for therapeutic reasons, it is clearly possible to treat certain injuries laparoscopically. Bleeding from minor injuries to the liver or the spleen can be controlled through the laparoscope (EL 4 [50, 53, 293]). Diaphragmatic lacerations (EL 4 [179, 248, 249]) and perforating stab wounds of the gastrointestinal tract can be sewn or stapled (EL 4 [53, 177, 293]). Nevertheless, the scarceness of clinical data prohibits a clear recommendation in favor of therapeutic laparoscopy for trauma.

Discussion

Available evidence clearly demonstrates the superiority of a laparoscopic approach in various emergency situations, but laparoscopy offers less or unclear benefit in other acute conditions. Therefore, a policy of laparoscopy for all patients with acute abdominal pain still seems unjustified, although laparoscopy will be to the advantage of the majority of patients. The initial usage of diagnostic procedures and imaging should aim to identify those patients who would probably not benefit from laparoscopy. On the other hand, it usually carries only minor disadvantages for a patient if a diagnostic laparoscopy has to be converted to an open procedure. Because the current guidelines deal with complex laparoscopic procedures, a low threshold toward early conversion is generally useful in order to avoid delays in the operating room.

Although the current recommendations address the most common diagnoses, some less prevalent causes of acute abdominal pain were not specifically discussed. Some of the more rare diagnoses were encountered in the cohort studies summarized in Table 1. These diseases include abdominal abscess, peritoneal tuberculosis, and intestinal volvulus. Due to their low occurrence, these diseases will probably never be studied in a randomized trial, but their relative importance in the treatment of an average patient is low. Laparoscopic therapy has been described to be useful for many of these conditions (EL 2b [265]; EL 4 [9, 127, 245]).

The panel also decided not to prepare separate recommendations on the usage of laparoscopy in children with acute abdominal pain. The disease spectrum of

pediatric acute abdominal pain is completely different compared to that of adults, but older children and adolescents are good candidates for laparoscopy (EL 4 [118, 287]). The value of specific procedures in pediatric surgery, such as laparoscopic appendectomy, is still under intensive debate. In consequence, these guidelines are valid only for adult patients. Also, there was no pediatric surgeon on the panel to define the possible role of laparoscopic surgery for pyloric stenosis, congenital malformations, and other disorders of the newborn or small child.

Future research should concentrate on those fields for which only low-level evidence is available. The current guidelines have identified some topics that have been described only in feasibility studies. It is highly desirable to supplement these studies with additional comparative data on effectiveness and cost-effectiveness. Because the EAES updates its guidelines regularly, such data are also important before stronger recommendations can be issued. On the other hand, in those fields for which there is good evidence, laparoscopic surgery has been shown to be highly beneficial. Therefore, optimism with regard to laparoscopy may prove to be justified. Laparoscopy has already had a major impact on the management of abdominal emergencies and has become an indispensable technique.

References

- Abbott J, Hawe J, Hunter D, Holmes M, Finn P, Garry R (2004) Laparoscopic excision of endometriosis: a randomized, placebo-controlled trial. *Fertil Steril* 82: 878–884
- Agresta F, De Simone P, Bedin N (2004) The laparoscopic approach in abdominal emergencies: a single-center 10-year experience. *J Soc Laparoendosc Surg* 8: 25–30
- Agresta F, Piazza A, Michelet I, Bedin N, Sartori CA (2000) Small bowel obstruction. Laparoscopic approach. *Surg Endosc* 14: 154–156
- Ahmad TA, Shelbaya E, Razek SA, Mohamed RA, Tajima Y, Ali SM, Sabet MM, Kanematsu T (2001) Experience of laparoscopic management in 100 patients with acute abdomen. *Hepatogastroenterology* 48: 733–736
- Ainsworth AP, Rafaelsen SR, Wamberg PA, Durup J, Pless TK, Mortensen MB (2003) Is there a difference in diagnostic accuracy and clinical impact between endoscopic ultrasonography and magnetic resonance cholangiopancreatography? *Endoscopy* 35: 1029–1032
- Alimoglu O, Ozkan OV, Sahin M, Akcakaya A, Eryilmaz R, Bas G (2003) Timing of cholecystectomy for acute biliary pancreatitis: outcomes of cholecystectomy on first admission and after current biliary pancreatitis. *World J Surg* 27: 256–259
- Allen LA, Schoon MG (1983) Laparoscopic diagnosis of acute pelvic inflammatory disease. *Br J Obstet Gynaecol* 90: 966–968
- Al-Mulhim AA (2000) Laparoscopic management of acute small bowel obstruction. Experience from a Saudi teaching hospital. *Surg Endosc* 14: 157–160
- Al-Mulhim AA (2004) Laparoscopic diagnosis of peritoneal tuberculosis. *Surg Endosc* 18: 1757–1761
- American Society of Colon and Rectal Surgeons (2000) Practice parameters for the treatment of sigmoid diverticulitis. *Dis Colon Rectum* 43: 289–297
- Anonymous (2002) NIH state-of-the-science statement on endoscopic retrograde cholangiopancreatography (ERCP) for diagnosis and therapy. *NIH Consens State Sci Statements* 19: 1–26
- Araujo-Teixeira JP, Rocha-Reis J, Costa-Cabral A, Barros H, Saraiva AC, Araujo-Teixeira AM (1999) Laparoscopia ou laparotomia dans la cholecystite aigue (200 cas). Comparaison des resultats et facteurs predisposant a la conversion. *Chirurgie* 124: 529–535
- Arguedas MR, Dupont AW, Wilcox CM (2001) Where do ERCP, endoscopic ultrasound, magnetic resonance cholangiopancreatography, and intraoperative cholangiography fit in the management of acute biliary pancreatitis? A decision analysis model. *Am J Gastroenterol* 96: 2892–2899
- Assalia A, Kopelman D, Hashmonai M (1997) Emergency minilaparotomy cholecystectomy for acute cholecystitis: prospective randomized trials—implications for the laparoscopic era. *World J Surg* 21: 534–539
- Avrutis O, Friedman SJ, Meshoum J, Haskel L, Adler S (2000) Safety and success of early laparoscopic cholecystectomy for acute cholecystitis. *Surg Laparosc Endosc Percutan Tech* 10: 200–207
- Ayub K, Imada R, Slavin J (2004) Endoscopic retrograde cholangiopancreatography in gallstone-associated acute pancreatitis. *Cochrane Database Syst Rev* CD003630
- Bailey IS, Rhodes M, O'Rourke N, Nathanson L, Fielding G (1998) Laparoscopic management of acute small bowel obstruction. *Br J Surg* 85: 84–87
- Bastug DF, Trammell SW, Boland JP, Mantz EP, Tiley EH 3rd (1991) Laparoscopic adhesiolysis for small bowel obstruction. *Surg Laparosc Endosc* 1: 259–262
- Bedirli A, Sözüer EM, Sakrak O, Babayigit H, Yilmaz Z (2003) Comparison of the results of early, delayed and elective surgery in biliary pancreatitis. *Turk J Gastroenterol* 14: 97–101
- Berber E, Engle KL, String A, Garland AM, Chang G, Macho J, Pearl JM, Siperstein AE (2000) Selective use of tube cholecystostomy with interval laparoscopic cholecystectomy in acute cholecystitis. *Arch Surg* 135: 341–346
- Bergamaschi R, Marvik R, Johnsen G, Thoresen JE, Ystgaard B, Myrvold HE (1999) Open vs laparoscopic repair of perforated peptic ulcer. *Surg Endosc* 13: 679–682
- Bevan CD, Johal BJ, Mumtaz G, Ridgway GL, Siddle NC (1995) Clinical, laparoscopic and microbiological findings in acute salpingitis; report on a United Kingdom cohort. *Br J Obstet Gynaecol* 102: 407–414
- Bhattacharya D, Ammori BJ (2005) Contemporary minimally invasive approaches to the management of acute cholecystitis: a review and appraisal. *Surg Laparosc Endosc Percutan Tech* 15: 1–8
- Bhattacharya D, Senapati PS, Hurle R, Ammori BJ (2002) Urgent versus interval laparoscopic cholecystectomy for acute cholecystitis: a comparative study. *J Hepatobiliary Pancreat Surg* 9: 538–542
- Bittner R, Leibl B, Kraft K, Butters M, Nick G, Ulrich M (1997) Laparoskopische Cholecystektomie in der Therapie der akuten Cholecystitis: Sofort- versus Intervalloperation. *Chirurg* 68: 237–243
- Bjorck M, Acosta S, Lindberg F, Troeng T, Bergqvist D (2002) Revascularization of the superior mesenteric artery after acute thromboembolic occlusion. *Br J Surg* 89: 923–927
- Boey J, Lee NW, Koo J, Lam PH, Wong J, Ong GB (1982) Immediate definitive surgery for perforated duodenal ulcers: a prospective controlled trial. *Ann Surg* 196: 338–344
- Boggi U, Di Candio G, Campatelli A, Oleggini M, Pietrabissa A, Filippini F, Bellini R, Mazzotta D, Mosca F (1999) Percutaneous cholecystostomy for acute cholecystitis in critically ill patients. *Hepatogastroenterology* 46: 121–125
- Borgstein PJ, Gordijn RV, Eijbsbouts QAJ, Cuesta MA (1997) Acute appendicitis—clear-cut case in men, a guessing game in young women. A prospective study on the role of laparoscopy. *Surg Endosc* 11: 923–927
- Borie F, Fingerhut A, Millat B (2003) Acute biliary pancreatitis, endoscopy, and laparoscopy. *Surg Endosc* 17: 1175–1180
- Borzellino G, de Manzoni G, Ricci F, Castaldini G, Guglielmi A, Cordiano C (1999) Emergency cholecystostomy and subsequent cholecystectomy for acute gallstone cholecystitis in the elderly. *Br J Surg* 86: 1521–1525
- Borzellino G, Tasselli S, Zerman G, Pedrazzani C, Manzoni G (2004) Laparoscopic approach to postoperative adhesive obstruction. *Surg Endosc* 18: 686–690
- Bove A, Bongarzone G, Serafini FM, Bonomo L, Dragani G, Palone F, Scotti U, Corbellini L (2004) L'approccio laparo-

- scopico alla colecistite acuta. Risultati e fattori predittivi di conversione. *G Chir* 25: 75–79
34. Bradley EL 3rd (1993) A clinically based classification system for acute pancreatitis. Summary of the International Symposium on Acute Pancreatitis, Atlanta, Ga, September 11 through 13, 1992. *Arch Surg* 128: 586–590
 35. British Society of Gastroenterology (1998) United Kingdom guidelines for the management of acute pancreatitis. *Gut* 42: S1–S13
 36. Brodsky A, Matter I, Sabo E, Cohen A, Abrahamson J, Eldar S (2000) Laparoscopic cholecystectomy for acute cholecystitis: can the need for conversion and the probability of complications be predicted? A prospective study. *Surg Endosc* 14: 755–760
 37. Bruwer F, Coetzer M, Warren BL (2003) Laparoscopic versus open surgical exploration in premenopausal women with suspected acute appendicitis. *S Afr J Surg* 41: 82–85
 38. Bryant TL, Umstot RK Jr (1996) Laparoscopic repair of an incarcerated obturator hernia. *Surg Endosc* 10: 437–438
 39. Budzynski A, Bobizynski A, Rembiasz K, Biesiada Z (2003) Laparoscopic emergency surgery. *Przegl Lek* 60: 20–24
 40. Byrne MF, Suhockl P, Mitchell RM, Pappas TN, Stiffler HL, Jowell PS, Branch MS, Baillie J (2003) Percutaneous cholecystostomy in patients with acute cholecystitis: experience of 45 patients at a U.S. referral center. *J Am Coll Surg* 197: 206–211
 41. Carbajo Caballero MA, Martín del Olmo JC, Blanco Álvarez JI, Cuesta de la Llave C, Atienza Sánchez R, Inglada Galiana L, Vaquero Puerta C (1998) Surgical treatment of the acute cholecystitis in the laparoscopic age. A comparative study: laparoscopy against laparotomy. *Rev Esp Enferm Dig* 90: 788–793
 42. Chak A, Hawes RH, Cooper GS, Hoffman B, Catalano MF, Wong RC, Herbener TE, Sivak MV Jr (1999) Prospective assessment of the utility of EUS in the evaluation of gallstone pancreatitis. *Gastrointest Endosc* 49: 599–604
 43. Champault G, Rizk N, Lauroy J, Olivares P, Belhassen A, Botutier P (1993) Douleurs iliaques droites de la femme. Approche diagnostique conventionnelle versus laparoscopie première. Étude contrôlée (65 cas). *Ann Chir* 47: 316–319
 44. Champault GG (1994) Laparoscopic treatment of perforated peptic ulcer. *Endosc Surg Allied Technol* 2: 117–118
 45. Chandler CF, Lane JS, Ferguson P, Thompson JE, Ashley SW (2000) Prospective evaluation of early versus delayed laparoscopic cholecystectomy for treatment of acute cholecystitis. *Am Surg* 66: 896–900
 46. Chang L, Lo S, Stabile BE, Lewis RJ, Toosie K, de Virgilio C (2000) Preoperative versus postoperative endoscopic retrograde cholangiopancreatography in mild to moderate gallstone pancreatitis: a prospective randomized trial. *Ann Surg* 231: 82–87
 47. Chang L, Moonka R, Stelzner M (2000) Percutaneous cholecystostomy for acute cholecystitis in veteran patients. *Am J Surg* 180: 198–202
 48. Chau CH, Tang CN, Siu WT, Ha JP, Li MK (2002) Laparoscopic cholecystectomy versus open cholecystectomy in elderly patients with acute cholecystitis: retrospective study. *Hong Kong Med J* 8: 394–399
 49. Cheema S, Brannigan AE, Johnson S, Delaney PV, Grace PA (2003) Timing of laparoscopic cholecystectomy in acute cholecystitis. *Ir J Med Sci* 172: 128–131
 50. Chen RJ, Fang JF, Lin BC, Hsu YB, Kao JL, Kao YC, Chen MF (1998) Selective application of laparoscopy and fibrin glue in the failure of nonoperative management of blunt hepatic trauma. *J Trauma* 44: 691–695
 51. Chiarugi M, Bucciante P, Decanini L, Balestri R, Lorenzetti L, Franceschi M, Cavina E (2001) “What you see is not what you get.” A plea to remove a “normal” appendix during diagnostic laparoscopy. *Acta Chir Belg* 101: 243–245
 52. Cho YP, Jung SM, Han MS, Jang HJ, Kim JS, Kim YH, Lee SG (2003) Role of diagnostic laparoscopy in managing acute mesenteric venous thrombosis. *Surg Laparosc Endosc Percutan Tech* 13: 215–217
 53. Chol YB, Lim KS (2003) Therapeutic laparoscopy for abdominal trauma. *Surg Endosc* 17: 421–427
 54. Chong AK, So JB, Ti TK (2001) Use of laparoscopy in the management of mesenteric venous thrombosis. *Surg Endosc* 15: 1042
 55. Chopra R, McVay C, Phillips E, Khalili TM (2003) Laparoscopic lysis of adhesions. *Am Surg* 69: 966–968
 56. Chosidow D, Johanet H, Montariol T, Kiehl R, Manceau C, Marmuse JP, Benhamou G (2000) Laparoscopy for acute small-bowel obstruction secondary to adhesions. *J Laparoendosc Adv Surg Tech A* 10: 155–159
 57. Chung RS, Diaz JJ, Chari V (1998) Efficacy of routine laparoscopy for the acute abdomen. *Surg Endosc* 12: 219–222
 58. Cloyd DW (1994) Laparoscopic repair of incarcerated paraesophageal hernias. *Surg Endosc* 8: 893–897
 59. Consensus Panel, Society for Surgery of the Alimentary Tract, American Gastroenterological Association, American Society for Gastrointestinal Endoscopy (2001) Management of the biliary tract in acute necrotizing pancreatitis. *J Gastrointest Surg* 5: 221–222
 60. Cougar P, Barrat C, Gayral F, et al. (2000) Société Française de Chirurgie Laparoscopique Le traitement laparoscopique de l'ulcère duodénal perforé. Résultats d'une étude rétrospective multicentrique. *Ann Chir* 125: 726–731
 61. Crofts TJ, Park KG, Steele RJ, Chung SS, Li AK (1989) A randomized trial of nonoperative treatment for perforated peptic ulcer. *N Engl J Med* 320: 970–973
 62. Cueto J, Diaz O, Garteiz D, Rodriguez M, Weber A (1997) The efficacy of laparoscopic surgery in the diagnosis and treatment of peritonitis. Experience with 107 cases in Mexico City. *Surg Endosc* 11: 366–370
 63. Cuschieri A, Hennessy TP, Stephens RB, Berci G (1988) Diagnosis of significant abdominal trauma after road traffic accidents: preliminary results of a multicentre clinical trial comparing minilaparoscopy with peritoneal lavage. *Ann R Coll Surg Engl* 70: 153–155
 64. Decadt B, Sussman L, Lewis MPN, Seeker A, Cohen L, Rogers C, Patel A, Rhodes M (1999) Randomized clinical trial of early laparoscopy in the management of acute non-specific abdominal pain. *Br J Surg* 86: 1383–1386
 65. Derveniz C, Johnson CD, Bassi C, Bradley E, Imrie CW, McMadlin MJ, Modlin I (1999) Diagnosis, objective assessment of severity, and management of acute pancreatitis. Santorini consensus conference. *Int J Pancreatol* 25: 195–210
 66. Druart ML, Van Hee R, Etienne J, et al. (1997) Laparoscopic repair of perforated duodenal ulcer. A prospective multicenter clinical trial. *Surg Endosc* 11: 1017–1020
 67. Easter DW, Cuschieri A, Nathanson LK, Lavelle-Jones M (1992) The utility of diagnostic laparoscopy for abdominal disorders. Audit of 120 patients. *Arch Surg* 127: 379–383
 68. Edwards MS, Cherr GS, Craven TE, Olsen AW, Plonk GW, Geary RL, Ligush JL, Hansen KJ (2003) Acute occlusive mesenteric ischemia: surgical management and outcomes. *Ann Vasc Surg* 17: 72–79
 69. Eldar S, Eitan A, Bickel A, Sabo E, Cohen A, Abrahamson J, Matter I (1999) The impact of patient delay and physician delay on the outcome of laparoscopic cholecystectomy for acute cholecystitis. *Am J Surg* 178: 303–307
 70. Eldar S, Sabo E, Nash E, Abrahamson J, Matter I (1997) Laparoscopic cholecystectomy for acute cholecystitis: prospective trial. *World J Surg* 21: 540–545
 71. Eldar S, Sabo E, Nash E, Abrahamson J, Matter I (1997) Laparoscopic versus open cholecystectomy in acute cholecystitis. *Surg Laparosc Endosc* 7: 407–414
 72. Eypasch E, Kum CK, Neugebauer E, et al. (2000) Laparoscopic appendectomy: EAES consensus development conference (1997) and updating comments (2000). In: Neugebauer E, Sauerland S, Troidl H (eds). *Recommendations for evidence-based surgery*. Springer, Paris, pp 15–23
 73. Eypasch E, Menningen R, Paul A, Troidl H (1993) Die Bedeutung der Laparoskopie bei der Diagnostik und Therapie des akuten Abdomens. *Zentralbl Chir* 118: 726–732
 74. Eypasch E, Spangenberg W, Ure B, Mennigen R, Troidl H (1994) Laparoskopische und konventionelle Übernähung perforierter peptischer Ulcera: eine Gegenüberstellung. *Chirurg* 65: 445–450
 75. Fagniez PL, Rotman N, Association de Recherche en Chirurgie (1998) Influence de la date de l'intervention chirurgicale sur le pronostic des pancréatites aiguës biliaires graves. *Chirurgie* 123: 368–372

76. Fan ST, Lai ECS, Mok FPT, Lo CML, Zheng SS, Wong J (1993) Early treatment of acute biliary pancreatitis by endoscopic papillotomy. *N Engl J Med* 328: 228–232
77. Faranda C, Barrat C, Catheline JM, Champault GG (2000) Two-stage laparoscopic management of generalized peritonitis due to perforated sigmoid diverticula; eighteen cases. *Surg Laparosc Endosc Percutan Tech* 10: 135–141
78. Feigel M, Thalmann C, Blessing H (1996) Laparoskopisch endoskopisch-präperitoneale kombinierte Hernien-Operation bei incarcerierter indirekter Inguinalhernie. *Chirurg* 67: 188–189
79. Fernandez H, Marchal L, Vincent Y (1998) Fertility after radical surgery for tubal pregnancy. *Fertil Steril* 70: 680–686
80. Ferrozzi L, Lippolis G, Petitti T, Carnevale D, Masi M (2004) La colecistectomia videolaparoscopica nella colecistite acuta: nostra esperienza. *G Chir* 25: 80–82
81. Ferzli G, Shapiro K, Chaudry G, Patel S (2004) Laparoscopic extraperitoneal approach to acutely incarcerated inguinal hernia. *Surg Endosc* 18: 228–231
82. Fevang BT, Fevang J, Lie SA, Soreide O, Svanes K, Viste A (2004) Long-term prognosis after operation for adhesive small bowel obstruction. *Ann Surg* 240: 193–201
83. Fevang BT, Fevang JM, Soreide O, Svanes K, Viste A (2003) Delay in operative treatment among patients with small bowel obstruction. *Scand J Surg* 92: 131–137
84. Fevang BT, Jensen D, Svanes K, Viste A (2002) Early operation or conservative management of patients with small bowel obstruction? *Eur J Surg* 168: 475–481
85. Fischer CP, Doherty D (2002) Laparoscopic approach to small bowel obstruction. *Semin Laparosc Surg* 9: 40–45
86. Flum DR, Morris A, Koepsell T, Dellinger EP (2001) Has misdiagnosis of appendicitis decreased over time? A population-based analysis. *J Am Med Assoc* 286: 1748–1753
87. Fölsch UR, Nitsche R, Ludtke R, Hilgers RA, Creutzfeldt W (1997) The German Study Group on Acute Biliary Pancreatitis Early ERCP and papillotomy compared with conservative treatment for acute biliary pancreatitis. *N Engl J Med* 336: 237–242
88. Franklin ME Jr, Dorman JP, Jacobs M, Plasencia G (1997) Is laparoscopic surgery applicable to complicated colonic diverticular disease? *Surg Endosc* 11: 1021–1025
89. Franklin ME Jr, Dorman JP, Pharand D (1994) Laparoscopic surgery in acute small bowel obstruction. *Surg Laparosc Endosc* 4: 289–296
90. Franklin ME Jr, Gonzalez JJ Jr, Miter DB, Glass JL, Paulson D (2004) Laparoscopic diagnosis and treatment of intestinal obstruction. *Surg Endosc* 18: 26–30
91. Gagner M (1996) Laparoscopic treatment of acute necrotizing pancreatitis. *Semin Laparosc Surg* 3: 21–28
92. Gaitán H, Angel E, Sánchez J, Gómez I, Sánchez L, Agudelo C (2002) Laparoscopic diagnosis of acute lower abdominal pain in women of reproductive age. *Int J Gynaecol Obstet* 76: 149–158
93. Garber SM, Korman J, Cosgrove JM, Cohen JR (1997) Early laparoscopic cholecystectomy for acute cholecystitis. *Surg Endosc* 11: 347–350
94. Geis WP, Kim HC (1995) Use of laparoscopy in the diagnosis and treatment of patients with surgical abdominal sepsis. *Surg Endosc* 9: 178–182
95. Gharaibeh KI, Qasaimeh GR, Al-Heiss H, Ammari F, Bani-Hani K, Al-Jaberi TM, Al-Natour S (2002) Effect of timing of surgery, type of inflammation, and sex on outcome of laparoscopic cholecystectomy for acute cholecystitis. *J Laparoendosc Adv Surg Tech A* 12: 193–198
96. Giger U, Michel JM, Vonlanthen R, Becker K, Kocher T, Krähenbühl L (2005) Laparoscopic cholecystectomy in acute cholecystitis: indication, technique, risk and outcome. *Langenbecks Arch Surg* 390: 373–380
97. Glavic Z, Begic L, Simlesa D, Rukavina A (2001) Treatment of acute cholecystitis. A comparison of open vs laparoscopic cholecystectomy. *Surg Endosc* 15: 398–401
98. Golash V, Willson PD (2005) Early laparoscopy as a routine procedure in the management of acute abdominal pain: a review of 1320 patients. *Surg Endosc* 19: 882–885
99. Graham A, Henley C, Mobley J (1991) Laparoscopic evaluation of acute abdominal pain. *J Laparoendosc Surg* 1: 165–168
100. Granlund A, Karlson BM, Elvin A, Rasmussen I (2001) Ultrasound-guided percutaneous cholecystostomy in high-risk surgical patients. *Langenbecks Arch Surg* 386: 212–217
101. Gray DT, Thorburn J, Lundorff P, Strandell A, Lindblom B (1995) A cost-effectiveness study of a randomised trial of laparoscopy versus laparotomy for ectopic pregnancy. *Lancet* 345: 1139–1143
102. Greenwald JA, McMullen HF, Coppa GF, Newman RM (2000) Standardization of surgeon-controlled variables: impact on outcome in patients with acute cholecystitis. *Ann Surg* 231: 339–344
103. Grunewald B, Keating J (1993) Should the “normal” appendix be removed at operation for appendicitis? *J R Coll Surg Edinb* 38: 158–160
104. Haan J, Kole K, Brunetti A, Kramer M, Scalea TM (2003) Nontherapeutic laparotomies revisited. *Am Surg* 69: 562–565
105. Habib FA, Kolachalam RB, Khilnani R, Preventza O, Mittal VK (2001) Role of laparoscopic cholel cystectomy in the management of gangrenous cholecystitis. *Am J Surg* 181: 71–75
106. Hallal AH, Amortegui JD, Jeroukhimov IM, Casillas J, Schulman CI, Manning RJ, Habib FA, Lo-pezz PP, Cohn SM, Sleeman D (2005) Magnetic resonance cholangiopancreatography accurately detects common bile duct stones in resolving gallstone pancreatitis. *J Am Coll Surg* 200: 869–875
107. Hamad GG, Broderick TJ (2000) Laparoscopic pancreatic necrosectomy. *J Laparoendosc Adv Surg Tech A* 10: 115–118
108. Hartwig W, Maksan SM, Foitzik T, Schmidt J, Herfarth C, Klar E (2002) Reduction in mortality with delayed surgical therapy of severe pancreatitis. *J Gastrointest Surg* 6: 481–487
109. Hatzidakis AA, Prassopoulos P, Petinarakis I, Sanidas E, Chrysos E, Chalkiaokakis G, Tsiftsis D, Gourtsoyannis NC (2002) Acute cholecystitis in high-risk patients: percutaneous cholecystostomy vs conservative treatment. *Eur Radiol* 12: 1778–1784
110. Heiss MM, Hüttl TP, Spelsberg FW, Lang RA, Jauch KW (2003) Möglichkeiten und Grenzen der laparoskopischen Magen-chirurgie. *Viszeralchirurgie* 38: 107–117
111. Ibrahim IM, Wolodiger F, Sussman B, Kahn M, Silvestri F, Sabar A (1996) laparoscopic management of acute small-bowel obstruction. *Surg Endosc* 10: 1012–1015
112. Irvin TT (1989) Abdominal pain: a surgical audit of 1190 emergency admissions. *Br J Surg* 76: 1121–1125
113. Ishihara T, Kubota K, Eda N, Ishibashi S, Haraguchi Y (1996) Laparoscopic approach to incarcerated inguinal hernia. *Surg Endosc* 10: 1111–1113
114. Isla A, Griniatsos J, Rodway A (2003) Single-stage definitive laparoscopic management in mild acute biliary pancreatitis. *J Laparoendosc Adv Surg Tech A* 13: 77–81
115. Ito K, Fujita N, Noda Y, Kobayashi G, Kimura K, Sugawara T, Horaguchi J (2004) Percutaneous cholecystostomy versus gall-bladder aspiration for acute cholecystitis: a prospective randomized controlled trial. *Am J Roentgenol* 183: 193–196
116. Jacobson TZ, Barlow DH, Garry R, Koninckx P (2001) Laparoscopic surgery for pelvic pain associated with endometriosis. *Cochrane Database Syst Rev* : CD001300
117. Jadallah FA, Abdul-Ghani AA, Tibblin S (1994) Diagnostic laparoscopy reduces unnecessary appendectomy in fertile women. *Eur J Surg* 160: 41–45
118. Jawad AJ, Al-Meshari A (1998) Laparoscopy for ovarian pathology in infancy and childhood. *Pediatr Surg Int* 14: 62–65
119. Johansson B, Hallerbäck B, Glise H, Johnsson E (1996) Laparoscopic suture closure of perforated peptic ulcer. A nonrandomized comparison with open surgery. *Surg Endosc* 10: 656–658
120. Johansson M, Thune A, Blomqvist A, Nelvin L, Lundell L (2003) Management of acute cholecystitis in the laparoscopic era: results of a prospective, randomized clinical trial. *J Gastrointest Surg* 7: 642–645
121. Johansson M, Thune A, Blomqvist A, Nelvin L, Lundell L (2004) Impact of choice of therapeutic strategy for acute cholecystitis on patient’s health-related quality of life: results of a randomized, controlled clinical trial. *Dig Surg* 21: 359–362
122. Johansson M, Thune A, Nelvin L, Stiernstam M, Westman B, Lundell L (2005) Randomized clinical trial of open versus laparoscopic cholecystectomy in the treatment of acute cholecystitis. *Br J Surg* 92: 44–49
123. Kaiwa Y, Namiki K, Matsumoto H (2003) Laparoscopic relief of reduction en masse of incarcerated inguinal hernia. *Surg Endosc* 17: 352

124. Kaltenthaler E, Vergel YB, Chilcott J, Thomas S, Blakeborough T, Walters SJ, Bouchier H (2004) A systematic review and economic evaluation of magnetic resonance cholangiopancreatography compared with diagnostic endoscopic retrograde cholangiopancreatography. *Health Technol Assess* 8:iii, 1–89
125. Kelly TR, Wagner DS (1988) Gallstone pancreatitis: a prospective randomized trial of the timing of surgery. *Surgery* 104: 600–605
126. Kim HJ, Lee SK, Kim MH, Yoo KS, Lim BC, Seo DW, Min YI (2000) Safety and usefulness of percutaneous transhepatic cholecystoscopy examination in high-risk surgical patients with acute cholecystitis. *Gastrointest Endosc* 52: 645–649
127. Kim SY, Moore JT (2003) Volvulus of the gallbladder: laparoscopic detorsion and removal. *Surg Endosc* 17: 1849
128. Kirby CP, Sparnon AL (2001) Active observation of children with possible appendicitis does not increase morbidity. *ANZ J Surg* 71: 412–413
129. Kirshtein B, Roy-Shapira A, Lantsberg L, Avinoach E, Mizrahi S (2005) Laparoscopic management of acute small bowel obstruction. *Surg Endosc*, 19: 464–467
130. Kirshtein B, Roy-Shapira A, Lantsberg L, Mandel S, Avinoach E, Mizrahi S (2003) The use of laparoscopy in abdominal emergencies. *Surg Endosc* 17: 1118–1124
131. Kiviluoto T, Sirén J, Luukkonen P, Kivilaakso E (1998) Randomised trial of laparoscopic versus open cholecystectomy for acute and gangrenous cholecystitis. *Lancet* 351: 321–325
132. Klein HM, Lensing R, Klosterhalfen B, Tons C, Gunther RW (1995) Diagnostic imaging of mesenteric infarction. *Radiology* 197: 79–82
133. Knight JS, Mercer SJ, Somers SS, Walters AM, Sadek SA, Toh SK (2004) Timing of urgent laparoscopic cholecystectomy does not influence conversion rate. *Br J Surg* 91: 601–604
134. Kohler L, Sauerland S, Neugebauer E, for the Scientific Committee of European Association Endoscopic Surgery (1999) Diagnosis and treatment of diverticular disease: results of a consensus development conference. *Surg Endosc* 13: 430–436
135. Kok KYY, Mathew VV, Yapp SKS (1999) Laparoscopic omental patch repair for perforated duodenal ulcer. *Am Surg* 65: 27–30
136. Kolla SB, Aggarwal S, Kumar A, Kumar R, Chumber S, Parshad R, Seenu V (2004) Early versus delayed laparoscopic cholecystectomy for acute cholecystitis: a prospective randomized trial. *Surg Endosc* 18: 1323–1327
137. Köninger J, Böttinger P, Redecke J, Butters M (2004) Laparoscopic repair of perforated gastroduodenal ulcer by running suture. *Langenbecks Arch Surg* 389: 11–16
138. Kontoravdis A, Chryssikopoulos A, Hassiakos D, Liapis A, Zourlas PA (1996) The diagnostic value of laparoscopy in 2365 patients with acute and chronic pelvic pain. *Int J Gynaecol Obstet* 52: 243–248
139. Koo KP, Thirlby RC (1996) Laparoscopic cholecystectomy in acute cholecystitis. What is the optimal timing for operation? *Arch Surg* 131: 540–545
140. Koperna T, Kissler M, Schulz F (1999) Laparoscopic versus open treatment of patients with acute cholecystitis. *Hepatogastroenterology* 46: 753–757
141. Kraemer M, Ohmann C, Leppert R, Yang Q (2000) Macroscopic assessment of the appendix at diagnostic laparoscopy is reliable. *Surg Endosc* 14: 625–633
142. Kronborg O (1993) Treatment of perforated sigmoid diverticulitis: a prospective randomized trial. *Br J Surg* 80: 505–507
143. Kum CK, Eypasch E, Lefering R, Paul A, Neugebauer E, Troldl H (1996) Laparoscopic cholecystectomy for acute cholecystitis: is it really safe? *World J Surg* 20: 43–49
144. Kum CK, Goh PMY, Isaac JR, Tekant Y, Ngoi SS (1994) Laparoscopic cholecystectomy for acute cholecystitis. *Br J Surg* 81: 1651–1654
145. Kuster GG, Domagk D (1996) Laparoscopic cholecystostomy with delayed cholecystectomy as an alternative to conversion to open procedure. *Surg Endosc* 10: 426–428
146. Lai PBS, Kwong KH, Leung KL, Kwok SPY, Chan ACW, Chung SCS, Lau WY (1998) Randomized trial of early versus delayed laparoscopic cholecystectomy for acute cholecystitis. *Br J Surg* 85: 764–767
147. Laine S, Rantala A, Gullichsen R, Ovaska J (1997) Laparoscopic appendectomy—is it worthwhile? A prospective, randomized study in young women. *Surg Endosc* 11: 95–97
148. Lam CM, Yuen AW, Chik B, Wai AC, Fan ST (2005) Laparoscopic surgery for common surgical emergencies: a population-based study. *Surg Endosc* 19: 774–779
149. Landau O, Kyzer S (2004) Emergent laparoscopic repair of incarcerated incisional and ventral hernia. *Surg Endosc* 18: 1374–1376
150. Lanzafame RJ (1993) Techniques for the simultaneous management of incarcerated ventral herniae and cholelithiasis via laparoscopy. *J Laparoendosc Surg* 3: 193–201
151. Larsson PG, Henriksson G, Olsson M, Boris J, Stroberg P, Tronstad SE, Skulman S (2001) Laparoscopy reduces unnecessary appendectomies and improves diagnosis in fertile women. A randomized study. *Surg Endosc* 15: 200–202
152. Lau H (2004) Laparoscopic repair of perforated peptic ulcer: a meta-analysis. *Surg Endosc* 18: 1013–1021
153. Lau WY, Leung KL, Kwong KH, Davey IC, Robertson C, Dawson JJ, Chung SC, Li AK (1996) A randomized study comparing laparoscopic versus open repair of perforated peptic ulcer using suture or sutureless technique. *Ann Surg* 224: 131–138
154. Lavonius MI, Ovaska J (2000) Laparoscopy in the evaluation of the incarcerated mass in groin hernia. *Surg Endosc* 14: 488–489
155. Lee FYJ, Leung KL, Lai PBS, Lau JWY (2001) Selection of patients for laparoscopic repair of perforated peptic ulcer. *Br J Surg* 88: 133–136
156. Lee HK, Han HS, Min SK, Lee JH (2005) Sex-based analysis of the outcome of laparoscopic cholecystectomy for acute cholecystitis. *Br J Surg* 92: 463–466
157. Lee KB, Wong SK (2002) Emergency laparoscopic surgery—the Changi General Hospital experience. *Ann Acad Med Singapore* 31: 155–157
158. Leibl BJ, Schmedt CG, Kraft K, Kraft B, Bittner R (2001) Laparoscopic transperitoneal hernia repair of incarcerated hernias: is it feasible? Results of a prospective study. *Surg Endosc* 15: 1179–1183
159. Leister I, Markus PM, Becker H (2003) Mesenteriale Ischämie: Hat die diagnostische Laparoskopie einen Stellenwert? *Chirurg* 74: 407–412
160. Leon EL, Metzger A, Tsiotos GG, Schlinkert RT, Sarr MG (1998) Laparoscopic management of small bowel obstruction: indications and outcome. *J Gastrointest Surg* 2: 132–140
161. Leppäniemi A, Haapiainen R (2003) Diagnostic laparoscopy in abdominal stab wounds: a prospective, randomized study. *J Trauma* 55: 636–645
162. Leppäniemi A, Salo J, Haapiainen R (1995) Complications of negative laparotomy for truncal stab wounds. *J Trauma* 38: 54–58
163. Levard H, Boudet MJ, Msika S, Molkhou JM, Hay JM, Laborde Y, Gillet M, Fingerhut A (2001) Laparoscopic treatment of acute small bowel obstruction: a multicentre retrospective study. *ANZ J Surg* 71: 641–646
164. Liao K, Ramirez J, Carryl S, Shaftan GW (1997) A new approach in the management of incarcerated hernia. Emergency laparoscopic hernia repair. *Surg Endosc* 11: 944–945
165. Lin E, Wear K, Tiszenkel HI (2002) Planned reduction of incarcerated groin hernias with hernia sac laparoscopy. *Surg Endosc* 16: 936–938
166. Liu CL, Lo CM, Chan JK, Poon RT, Lam CM, Fan ST, Wong J (2001) Detection of choledocholithiasis by EUS in acute pancreatitis: a prospective evaluation in 100 consecutive patients. *Gastrointest Endosc* 54: 325–330
167. Liu S, Chen F, Chen H (1995) An evaluation of nonsurgical treatment for severe acute pancreatitis. *Zhonghua Wai Ke Za Zhi* 33: 545–547
168. Lo CM, Fan ST, Liu CL, Lai EC, Wong J (1997) Early decision for conversion of laparoscopic to open cholecystectomy for treatment of acute cholecystitis. *Am J Surg* 173: 513–517
169. Lo CM, Liu CL, Fan ST, Lai EC, Wong J (1998) Prospective randomized study of early versus delayed laparoscopic cholecystectomy for acute cholecystitis. *Ann Surg* 227: 461–467
170. Lujan JA, Parrilla P, Robles R, Marin P, Torralba JA, Garcia-Ayllon J (1998) Laparoscopic cholecystectomy vs open cholecystectomy in the treatment of acute cholecystitis: a prospective study. *Arch Surg* 133: 173–175

171. Lunderoff P, Thorburn J, Hahlin M, Kallfelt B, Lindblom B (1991) Laparoscopic surgery in ectopic pregnancy. A randomized trial versus laparotomy. *Acta Obstet Gynecol Scand* 70: 343–348
172. Lunderoff P, Thorburn J, Lindblom B (1992) Fertility outcome after conservative surgical treatment of ectopic pregnancy evaluated in a randomized trial. *Fertil Steril* 57: 998–1002
173. Madan AK, Aliabadi-Wahle S, Tesi D, Flint LM, Steinberg SM (2002) How early is early laparoscopic treatment of acute cholecystitis? *Am J Surg* 183: 232–236
174. Magos AL, Baumann R, Turnbull AC (1989) Managing gynaecological emergencies with laparoscopy. *Br Med J* 299: 371–374
175. Mais V, Ajossa S, Piras B, Marongiu D, Guerriero S, Melis GB (1995) Treatment of nonendometriotic benign adnexal cysts; a randomized comparison of laparoscopy and laparotomy. *Obstet Gynecol* 86: 770–774
176. Majewski WD (2005) Long-term outcome, adhesions, and quality of life after laparoscopic open surgical therapies for acute abdomen: follow-up of a prospective trial. *Surg Endosc* 19: 81–90
177. Mathonnet M, Peyrou P, Gainant A, Bouvier S, Cubertafond P (2003) Role of laparoscopy in blunt perforations of the small bowel. *Surg Endosc* 17: 641–645
178. Matsuda M, Nishiyama M, Hanai T, Saeki S, Watanabe T (1995) Laparoscopic omental patch repair for perforated peptic ulcer. *Ann Surg* 221: 236–240
179. Matthews BD, Bui H, Harold KL, Kercher KW, Adrales G, Park A, Sing RF, Heniford BT (2003) Laparoscopic repair of traumatic diaphragmatic injuries. *Surg Endosc* 17: 254–258
180. Mehendale VG, Shenoy SN, Joshi AM, Chaudhari NC (2002) Laparoscopic versus open surgical closure of perforated duodenal ulcers: a comparative study. *Indian J Gastroenterol* 21: 222–224
181. Merriam LT, Kanaan SA, Dawes LG, Angelos P, Prystowsky JB, Rege RV, Joehl RJ (1999) Gangrenous cholecystitis: analysis of risk factors and experience with laparoscopic cholecystectomy. *Surgery* 126: 680–686
182. Mier J, Luque-de León E, Castillo A, Robledo F, Blanco R (1997) Early versus late necrosectomy in severe necrotizing pancreatitis. *Am J Surg* 173: 71–75
183. Mikkelsen AL, Felding C (1990) Laparoscopy and ultrasound examination in women with acute pelvic pain. *Gynecol Obstet Invest* 30: 162–164
184. Millat B, Fingerhut A, Borie F (2000) Surgical treatment of complicated duodenal ulcers: controlled trials. *World J Surg* 24: 299–306
185. Miserez M, Eypasch E, Spangenberger W, Lefering R, Troidl H (1996) Laparoscopic and conventional closure of perforated peptic ulcer. A comparison. *Surg Endosc* 10: 831–836
186. Moberg AC, Berndsen F, Palmquist I, Petersson U, Resch T, Montgomery A (2005) Randomized clinical trial of laparoscopic versus open appendicectomy for confirmed appendicitis. *Br J Surg* 92: 298–304
187. Morino M, Castagna E, Rosso E, Mao P (2004) Non-specific acute abdominal pain: a randomized controlled trial comparing early laparoscopy vs. observation [abstract]. *Surg Endosc* 18: S2
188. Mouret P, Francois Y, Vignal J, Barth X, Lombard-Platet R (1990) Laparoscopic treatment of perforated peptic ulcer. *Br J Surg* 77: 1006
189. Murphy AA, Nager CW, Wujek JJ, Kettel LM, Torp VA, Chin HG (1992) Operative laparoscopy versus laparotomy for the management of ectopic pregnancy: a prospective trial. *Fertil Steril* 57: 1180–1185
190. Murphy MK, Black NA, Lamping DL, McKee CM, Sanderson CFB, Askham J, Marteau T (1998) Consensus development methods, and their use in clinical guideline development. *Health Technol Assessment* 2: i–iv, 1–88
191. Næsgaard JM, Edwin B, Reiertsens O, Trondsen E, Færden AE, Rosseland AR (1999) Laparoscopic and open operation in patients with perforated peptic ulcer. *Eur J Surg* 165: 209–214
192. Nagle A, Ujiki M, Denham W, Murayama K (2004) Laparoscopic adhesiolysis for small bowel obstruction. *Am J Surg* 187: 464–470
193. Nagy AG, James D (1989) Diagnostic laparoscopy. *Am J Surg* 157: 490–493
194. Nathanson LK, Easter DW, Cuschieri A (1990) Laparoscopic repair peritoneal toilet of perforated duodenal ulcer. *Surg Endosc* 4: 232–233
195. Nathens AB, Gurtis JR, Beale RJ, Cook DJ, Moreno RP, Romand JA, Skerrett SJ, Stapleton RD, Ware LB, Waldmann CS (2004) Management of the critically ill patient with severe acute pancreatitis. *Crit Care Med* 32: 2524–2536
196. Navarrete Aulestia S, Cantele H, Leyba JL, Navarrete M, Navarrete Llopla S (2003) Laparoscopic diagnosis and treatment in gynecologic emergencies. *J Soc Laparoendosc Surg* 7: 239–242
197. Navez B, Arimont JM, Guiot P (1998) Laparoscopic approach in acute small bowel obstruction. A review of 68 patients. *Hepato-gastroenterology* 45: 2146–2150
198. Navez B, d’Udekem Y, Cambier E, Richir C, de Pierpont B, Guiot P (1995) Laparoscopy for management of nontraumatic acute abdomen. *World J Surg* 19: 382–386
199. Navez B, Mutter D, Russier Y, Vix M, Jamali F, Lipski D, Cambier E, Guiot P, Leroy J, Marescaux J (2001) Safety of laparoscopic approach for acute cholecystitis: retrospective study of 609 cases. *World J Surg* 25: 1352–1356
200. Neoptolemos JP, Carr-Locke DL, London NJ, Bailey IA, James D, Fossard DP (1988) Controlled trial of urgent endoscopic retrograde cholangiopancreatography and endoscopic sphincterotomy versus conservative treatment for acute pancreatitis due to gallstones. *Lancet* 2: 979–983
201. Neoptolemos JP, Shaw DE, Carr-Locke DL (1989) A multivariate analysis of preoperative risk factors in patients with common bile duct stones. Implications for treatment *Ann Surg* 209: 157–161
202. Nicolau AE, Ionescu G, Iordache F, Mehic R, Spataru A (2002) Sutura laparoscopica sau sutura deschisa in ulcerul duodenal recent perforat la tineri. *Chirurgia (Bucur)* 97: 305–311
203. Nitke S, Goldman GA, Fisch B, Kaplan B, Ovadia J (1996) The management of dermoid cysts—a comparative study of laparoscopy and laparotomy. *Isr J Med Sci* 32: 1177–1179
204. Nordback I, Sisto T (1991) Ultrasonography and computed tomography in the diagnosis of portomesenteric vein thrombosis. *Int Surg* 76: 179–182
205. Olsen JB, Myrén CJ, Haahr PE (1993) Randomized study of the value of laparoscopy before appendicectomy. *Br J Surg* 80: 922–923
206. O’Sullivan GC, Murphy D, O’Brien MG, Ireland A (1996) Laparoscopic management of generalized peritonitis due to perforated colonic diverticula. *Am J Surg* 171: 432–434
207. Ou CS, Rowbotham R (2000) Laparoscopic diagnosis and treatment of nontraumatic acute abdominal pain in women. *J Laparoendosc Adv Surg Tech A* 10: 41–45
208. Paimela H, Oksala NK, Kivilaakso E (2004) Surgery for peptic ulcer today. A study on the incidence, methods and mortality in surgery for peptic ulcer in Finland between 1987 and 1999. *Dig Surg* 21: 185–191
209. Pamoukian VN, Gagner M (2001) Laparoscopic necrosectomy for acute necrotizing pancreatitis. *J Hepatobiliary Pancreat Surg* 8: 221–223
210. Papi C, Catarci M, D’Ambrosio L, Gili L, Koch M, Grassi GB, Capurso L (2004) Timing of cholecystectomy for acute calculous cholecystitis: a meta-analysis. *Am J Gastroenterol* 99: 147–155
211. Paterson-Brown S, Eckersley JRT, Sim AJW, Dudley HAF (1986) Laparoscopy as an adjunct to decision making in the “acute abdomen”. *Br J Surg* 73: 1022–1024
212. Patient Care Committee of the Society for Surgery Alimentary Tract (1999) Surgical treatment of diverticulitis. *J Gastrointest Surg* 3: 212–213
213. Patterson EJ, McLoughlin RF, Mathieson JR, Cooperberg PL, MacFarlane JK (1996) An alternative approach to acute cholecystitis. Percutaneous cholecystostomy and interval laparoscopic cholecystectomy. *Surg Endosc* 10: 1185–1188
214. Paul A, Millat B, Holthausen U, Sauerland S, Neugebauer E, for the Scientific Committee of European Association for Endoscopic Surgery (1998) Diagnosis and treatment of common bile duct stones (CBDS). Results of a consensus development conference. *Surg Endosc* 12: 856–864
215. Peng WK, Sheikh Z, Nixon SJ, Paterson-Brown S (2005) Role of laparoscopic cholecystectomy in the early management of acute gallbladder disease. *Br J Surg* 92: 586–591

216. Perko MJ (2001) Duplex ultrasound for assessment of superior mesenteric artery blood flow. *Eur J Vasc Endovasc Surg* 21: 106–117
217. Pessaux P, Lebigot J, Tuech JJ, Regenet N, Aubé C, Ridereau C, Arnaud JP (2000) Cholecystostomie percutanée dans les cholécystites aiguës chez les patients à haut risque. *Ann Chir* 125: 738–743
218. Pessaux P, Regenet N, Tuech JJ, Rouge C, Bergamaschi R, Arnaud JP (2001) Laparoscopic versus open cholecystectomy: a prospective comparative study in the elderly with acute cholecystitis. *Surg Laparosc Endosc Percutan Tech* 11: 252–255
219. Pessaux P, Tuech JJ, Derouet N, Rouge C, Regenet N, Arnaud JP (2000) Laparoscopic cholecystectomy in the elderly: a prospective study. *Surg Endosc* 14: 1067–1069
220. Pessaux P, Tuech JJ, Rouge C, Duplessis R, Cervi C, Arnaud JP (2000) Laparoscopic cholecystectomy in acute cholecystitis. A prospective comparative study in patients with acute vs. chronic cholecystitis. *Surg Endosc* 14: 358–361
221. Prat F, Edery J, Meduri B, et al. (2001) Early EUS of the bile duct before endoscopic sphincterotomy for acute biliary pancreatitis. *Gastrointest Endosc* 54: 724–729
222. Promecene PA (2002) Laparoscopy in gynecologic emergencies. *Semin Laparosc Surg* 9: 64–75
223. Pugliese R, Di Lernia S, Sansonna F, Scandroglia I, Maggioni D, Ferrari C, Costanzi A, Chiara O (2004) Laparoscopic treatment of sigmoid diverticulitis: a retrospective review of 103 cases. *Surg Endosc* 18: 1344–1348
224. Rau HG, Meyer G, Maiwald G, Schardey M, Merkle R, Lange V, Schildberg FW (1994) Konventionelle oder laparoskopische Cholecystektomie zur Behandlung der akuten Cholecystitis? *Chirurg* 65: 1121–1125
225. Reiertsen O, Rosseland AR, Hoivik B, Solheim K (1985) Laparoscopy in patients admitted for acute abdominal pain. *Acta Chir Scand* 151: 52–524
226. Renz BM, Feliciano DV (1995) Unnecessary laparotomies for trauma: a prospective study of morbidity. *J Trauma* 38: 350–356
227. Rhodes M, Sussman L, Cohen L, Lewis MP (1998) Randomised trial of laparoscopic exploration of common bile duct versus postoperative endoscopic retrograde cholangiography for common bile duct stones. *Lancet* 351: 159–161
228. Ricci F, Castaldini G, de Manzoni G, Botzellino G, Rodella L, Kind R (1997) Minimally invasive treatment of acute biliary pancreatitis. *Surg Endosc* 11: 1179–1182
229. Romagnuolo J, Currie G (2005) Noninvasive vs. selective invasive biliary imaging for acute biliary pancreatitis: an economic evaluation by using decision tree analysis. *Gastrointest Endosc* 61: 86–87
230. Rutledge D, Jones D, Reqe R (2000) Consequences of delay in surgical treatment of biliary disease. *Am J Surg* 180: 466–469
231. Salky BA, Edye MB (1998) The role of laparoscopy in the diagnosis and treatment of abdominal pain syndromes. *Surg Endosc* 12: 911–914
232. Sanna A, Adani GL, Anania G, Donini A (2003) The role of laparoscopy in patients with suspected peritonitis: experience of a single institution. *J Laparoendosc Adv Surg Tech A* 13: 17–19
233. Sauerland S, Lefering R, Neugebauer E (2004) Laparoscopic versus open surgery for suspected appendicitis [Cochrane Review]. In: the Cochrane Collaboration (ed). *The cochrane database of systematic reviews, Vol. IV/2004 [CD-ROM]*. Oxford Update Software
234. Scheiman JM, Carlos RC, Barnett JL, Elta GH, Nostrant TT, Chey WD, Francis IR, Nandi PS (2001) Can endoscopic ultrasound or magnetic resonance cholangiopancreatography replace ERCP in patients with suspected biliary disease? A prospective trial and cost analysis. *Am J Gastroenterol* 96: 2900–2904
235. Schlachta CM, Mamazza J, Poulin EC (1999) Laparoscopic sigmoid resection for acute and chronic diverticulitis. An outcomes comparison with laparoscopic resection for nondiverticular disease. *Surg Endosc* 13: 649–653
236. Schrenk P, Woisetschlager R, Wayand WU, Rieger R, Sulzbacher H (1994) Diagnostic laparoscopy: a survey of 92 patients. *Am J Surg* 168: 348–351
237. Schroder T, Sainio V, Kivisaari L, Puolakkainen P, Kivilaakso E, Lempinen M (1991) Pancreatic resection versus peritoneal lavage in acute necrotizing pancreatitis. A prospective randomized trial. *Ann Surg* 214: 663–666
238. Schwandner O, Farke S, Fischer F, Eckmann C, Schiedeck TH, Bruch HP (2004) Laparoscopic colectomy for recurrent and complicated diverticulitis: a prospective study of 396 patients. *Langenbecks Arch Surg* 389: 97–103
239. Scierski A (2004) Laparoscopic operations of incarcerated inguinal and femoral hernias. *Wiad Lek* 57: 245–248
240. Seelig MH, Seelig SK, Behr C, Schonleben K (2003) Comparison between open and laparoscopic technique in the management of perforated gastroduodenal ulcers. *J Clin Gastroenterol* 37: 226–229
241. Senapati PS, Bhattacharya D, Harinath G, Ammori BJ (2003) A survey of the timing and approach to the surgical management of cholelithiasis in patients with acute biliary pancreatitis and acute cholecystitis in the UK. *Ann R Coll Surg Engl* 85: 306–312
242. Serralta AS, Bueno JL, Planells MR, Rodero DR (2003) Prospective evaluation of emergency versus delayed laparoscopic cholecystectomy for early cholecystitis. *Surg Laparosc Endosc Percutan Tech* 13: 71–75
243. Shalaby R, Desoky A (2001) Laparoscopic approach to small intestinal obstruction in children: a preliminary experience. *Surg Laparosc Endosc Percutan Tech* 11: 301–305
244. Siu WT, Chau CH, Law BK, Tang CN, Ha PY, Li MK (2004) Routine use of laparoscopic repair for perforated peptic ulcer. *Br J Surg* 91: 481–484
245. Siu WT, Law BK, Tang CN, Chau CH, Li MK (2003) Laparoscopic management of omental torsion secondary to an occult inguinal hernia. *J Laparoendosc Adv Surg Tech A* 13: 199–201
246. Siu WT, Leong HT, Law BKB, Chau CH, Li ACN, Fung KH, Tai YP, Li MKW (2002) Laparoscopic repair for perforated peptic ulcer: a randomized controlled trial. *Ann Surg* 235: 313–319
247. Siu WT, Leong HT, Li MK (1997) Single stitch laparoscopic omental patch repair of perforated peptic ulcer. *J R Coll Surg Edinb* 42: 92–94
248. Smith CH, Novick TL, Jacobs DG, Thomason MH (2000) Laparoscopic repair of a ruptured diaphragm secondary to blunt trauma. *Surg Endosc* 14: 501–502
249. Smith RS, Fry WR, Morabito DJ, Koehler RH, Organ CH Jr (1995) Therapeutic laparoscopy in trauma. *Am J Surg* 170: 632–637
250. Sø JBY, Kum CK, Fernanes ML, Goh P (1996) Comparison between laparoscopic and conventional omental patch repair for perforated duodenal ulcer. *Surg Endosc* 10: 1060–1063
251. Soper DE (1991) Diagnosis and laparoscopic grading of acute salpingitis. *Am J Obstet Gynecol* 164: 1370–1376
252. Sözüer EM, Bedirli A, Ulusal M, Kayhan E, Yilmaz Z (2000) Laparoscopy for diagnosis and treatment of acute abdominal pain. *J Laparoendosc Adv Surg Tech A* 10: 203–207
253. Sreenarasimhaiah J (2003) Diagnosis and management of intestinal ischaemic disorders. *Br Med J* 326: 1372–1376
254. Strickland P, Lourie DJ, Suddleson EA, Blitz JB, Stain SC (1999) Is laparoscopy safe and effective for treatment of acute small-bowel obstruction? *Surg Endosc* 13: 695–698
255. Suc B, Escat J, Cherqui P, Fourtanier G, Hay JM, Fingerhut A, Millat B (1998) French Associations for Surgical Research Surgery endoscopy as primary treatment in symptomatic patients with suspected common bile duct stones: a multicenter randomized trial. *Arch Surg* 133: 702–708
256. Sugarbaker PH, Sanders JH, Bloom BS, Wilson RE (1975) Preoperative laparoscopy in diagnosis of acute abdominal pain. *Lancet* 1: 442–445
257. Sugiyama M, Tokuhara M, Atomi Y (1998) Is percutaneous cholecystostomy the optimal treatment for acute cholecystitis in the very elderly? *World J Surg* 22: 459–463
258. Suter M, Meyer A (2001) A 10-year experience with the use of laparoscopic cholecystectomy for acute cholecystitis: is it safe? *Surg Endosc* 15: 1187–1192
259. Suter M, Zermatten P, Halkic N, Martinet O, Bettschart V (2000) Laparoscopic management of mechanical small bowel obstruction: are there predictors of success or failure? *Surg Endosc* 14: 478–483
260. Sutton CJG, Pooley AS, Ewen SP, Haines P (1997) Follow-up report on a randomized controlled trial of laser laparoscopy in

- the treatment of pelvic pain associated with minimal to moderate endometriosis. *Fertil Steril* 68: 1070–1074
261. Suzuki K, Umehara Y, Kimura T (2003) Elective laparoscopy for small bowel obstruction. *Surg Laparosc Endosc Percutan Tech* 13: 254–256
 262. Swank DJ, Swank-Bordewijk SC, Hop WC, van Erp WF, Janssen IM, Bonjer HJ, Jeekel J (2003) Laparoscopic adhesiolysis in patients with chronic abdominal pain: a blinded randomised controlled multi-centre trial. *Lancet* 361: 1247–1251
 263. Targarona EM, Balague C, Espert JJ, Perez Ayuso RM, Ros E, Navarro S, Bordas J, Teres J, Trias M (1995) Laparoscopic treatment of acute biliary pancreatitis. *Int Surg* 80: 365–368
 264. Taylor EW, Kennedy CA, Dunham RH, Bloch JH (1995) Diagnostic laparoscopy in women with acute abdominal pain. *Surg Laparosc Endosc* 5: 125–128
 265. Teague WJ, Ackroyd R, Watson DI, Devitt PG (2000) Changing patterns in the management of gastric volvulus over 14 years. *Br J Surg* 87: 353–361
 266. Teh SH, O'Ceallaigh S, McKeon JG, O'Donohue MK, Tanner WA, Keane FB (2000) Should an appendix that look "normal" be removed at diagnostic laparoscopy for acute right iliac fossa pain? *Eur J Surg* 166: 388–389
 267. Teisala K, Heinonen PK, Punnonen R (1990) Laparoscopic diagnosis and treatment of acute pyosalpinx. *J Reprod Med* 35: 19–21
 268. Teixeira JP, Saraiva AC, Cabral AC, Barros H, Reis JR, Teixeira A (2000) Conversion factors in laparoscopic cholecystectomy for acute cholecystitis. *Hepatogastroenterology* 47: 626–630
 269. Trias M, Targarona EM, Ros E, Bordas JM, Perez Ayuso RM, Balague C, Pros I, Teres J (1997) Prospective evaluation of a minimally invasive approach for treatment of bile-duct calculi in the high-risk patient. *Surg Endosc* 11: 632–635
 270. Trowbridge RL, Rutkowski NK, Shojania KG (2003) Does this patient have acute cholecystitis? *J Am Med Assoc* 289: 80–86
 271. Tschudi J, Wagner M, Klaiber C (1993) Laparoskopische Operation einer incarcerierten Obturatorhernie mit assistierter Darmresektion. *Chirurg* 64: 827–828
 272. Tsumura H, Ichikawa T, Hiyama E, Murakami Y (2004) Laparoscopic and open approach in perforated peptic ulcer. *Hepatogastroenterology* 51: 1536–1539
 273. Uchiyama K, Onishi H, Tani M, Kinoshita H, Ueno M, Yamaue H (2004) Timing of laparoscopic cholecystectomy for acute cholecystitis with cholelithiasis. *Hepatogastroenterology* 51: 346–348
 274. Uhl W, Müller CA, Krähenbühl L, Schmid SW, Scholzel S, Büchler MW (1999) Acute gallstone pancreatitis: timing of laparoscopic cholecystectomy in mild and severe disease. *Surg Endosc* 13: 1070–1076
 275. UK Working Party on Acute Pancreatitis (2005) UK guidelines for the management of acute pancreatitis. *Gut* 54: iii1–iii9
 276. Urbach DR, Khajanchee YS, Jobe BA, Standage BA, Hansen PD, Swanstrom LL (2001) Cost-effective management of common bile duct stones: a decision analysis of the use of endoscopic retrograde cholangiopancreatography (ERCP), intraoperative cholangiography, and laparoscopic bile duct exploration. *Surg Endosc* 15: 4–13
 277. van Dalen R, Bagshaw PF, Dobbs BR, Robertson GM, Lynch AC, Frizelle FA (2003) The utility of laparoscopy in the diagnosis of acute appendicitis in women of reproductive age. *Surg Endosc* 17: 1311–1313
 278. van den Broek WT, Bijnen AB, de Ruiter P, Gouma DJ (2001) A normal appendix found during diagnostic laparoscopy should not be removed. *Br J Surg* 88: 251–254
 279. Vermesh M, Silva PD, Rosen GF, Stein AL, Fossum GT, Sauer MV (1989) Management of unruptured ectopic gestation by linear salpingostomy: a prospective, randomized clinical trial of laparoscopy versus laparotomy. *Obstet Gynecol* 73: 400–404
 280. Vetrhus M, Soreide O, Nesvik I, Sondenaa K (2003) Acute cholecystitis: delayed surgery or observation. A randomized clinical trial. *Scand J Gastroenterol* 38: 985–990
 281. Villavicencio RT, Aucar JA (1999) Analysis of laparoscopy in trauma. *J Am Coll Surg* 189: 11–20
 282. Waclawiczek HW, Schneeberger V, Bekk A, Dinnewitzer A, Sungler P, Boeckl O (1997) Der Stellenwert der diagnostischen Laparoskopie und minimal-invasiver Verfahren beim akuten Abdomen. *Zentralbl Chir* 122: 1108–1112
 283. Watson SD, Saye W, Hollier PA (1993) Combined laparoscopic incarcerated herniorrhaphy and small bowel resection. *Surg Laparosc Endosc* 3: 106–108
 284. Williams SB, Greenspon J, Young HA, Orkin BA (2005) Small bowel obstruction: conservative vs. surgical management. *Dis Colon Rectum* 48: 1140–1146
 285. Willsher PC, Sanabria JR, Gallinger S, Rossi L, Strasberg S, Litwin DE (1999) Early laparoscopic cholecystectomy for acute cholecystitis: a safe procedure. *J Gastrointest Surg* 3: 50–53
 286. Wilson RG, Macintyre IM, Nixon SJ, Saunders JH, Varma JS, King PM (1992) Laparoscopic cholecystectomy as a safe and effective treatment for severe acute cholecystitis. *Br Med J* 305: 394–396
 287. Wolfman WL, Kreutner K (1984) Laparoscopy in children and adolescents. *J Adolesc Health Care* 5: 261–265
 288. Wong SK, Yu SC, Lam YH, Chung SS (1999) Percutaneous cholecystostomy and endoscopic cholelithotripsy in the management of acute cholecystitis. *Surg Endosc* 13: 48–52
 289. Wullstein C, Gross E (2003) Laparoscopic compared with conventional treatment of acute adhesive small bowel obstruction. *Br J Surg* 90: 1147–1151
 290. Yau KK, Siu WT, Chau CH, Yang PC, Li MK (2005) Laparoscopic management of incarcerated obturator hernia. *Can J Surg* 48: 76–77
 291. Yuen PM, Yu KM, Yip SK, Lau WC, Rogers MS, Chang A (1997) A randomized prospective study of laparoscopy and laparotomy in the management of benign ovarian masses. *Am J Obstet Gynecol* 177: 109–114
 292. Zamir G, Reissman P (1998) Diagnostic laparoscopy in mesenteric ischemia. *Surg Endosc* 12: 390–393
 293. Zantut LF, Ivatury RR, Smith RS, Kawahara NT, Porter JM, Fry WR, Poggetti R, Birolini D, Organ CH Jr (1997) Diagnostic and therapeutic laparoscopy for penetrating abdominal trauma: a multicenter experience. *J Trauma* 42: 825–831
 294. Zeitoun G, Laurent A, Rouffet F, Hay J, Fingerhut A, Paquet J, Peillon C de Association Recher che en Chirurgie (2000) Multi-centre, randomized clinical trial of primary versus secondary moid resection in generalized peritonitis complicating sigmoid diverticulitis. *Br J Surg* 87: 1366–1374
 295. Z'Graggen K, Metzger A, Birrer S, Klaiber C (1995) Die laparoskopische Cholecystektomie als Standardtherapie bei der akuten Cholecystitis. Eine prospektive Studie. *Chirurg* 66: 366–370
 296. Zhang J, Duan ZQ, Song QB, Luo YW, Xin SJ, Zhang Q (2004) Acute mesenteric venous thrombosis: a better outcome achieved through improved imaging techniques and a changed policy of clinical management. *Eur J Vase Endovasc Surg* 28: 329–334
 297. Zhu JF, Fan XH, Zhang XH (2001) Laparoscopic treatment of severe acute pancreatitis. *Surg Endosc* 15: 146–148
 298. Zittel TT, Jehle EC, Becker HD (2000) Surgical management of peptic ulcer disease today—indication, technique and outcome. *Langenbecks Arch Surg* 385: 84–96